

Research

Open Access

## The activation of nicotinic acetylcholine receptors enhances the inhibitory synaptic transmission in the deep dorsal horn neurons of the adult rat spinal cord

Daisuke Takeda<sup>\*1,2</sup>, Terumasa Nakatsuka<sup>3</sup>, Jianguo G Gu<sup>4</sup> and Munehito Yoshida<sup>1</sup>

Address: <sup>1</sup>Department of Orthopaedic Surgery, Wakayama Medical University, 811-1 Kimiidera, Wakayama 641-8510, Japan, <sup>2</sup>Department of Physiology, Kansai University of Health Sciences, Osaka 590-0482, Japan, <sup>3</sup>Department of Physiology, Faculty of Medicine, Saga University, Saga 849-8501, Japan and <sup>4</sup>Brain Institute and Department of Oral Surgery, Division of Neuroscience, College of Dentistry, University of Florida, Gainesville, Florida 32610, USA

Email: Daisuke Takeda<sup>\*</sup> - dtakeda@kansai.ac.jp; Terumasa Nakatsuka - nakatsuk@cc.saga-u.ac.jp; Jianguo G Gu - jgu@dental.ufl.edu; Munehito Yoshida - myoshida@wakayama-med.ac.jp

<sup>\*</sup> Corresponding author

Published: 25 September 2007

Received: 17 July 2007

*Molecular Pain* 2007, **3**:26 doi:10.1186/1744-8069-3-26

Accepted: 25 September 2007

This article is available from: <http://www.molecularpain.com/content/3/1/26>

© 2007 Takeda et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

Somatosensory information can be modulated by nicotinic acetylcholine receptors (nAChRs) in the superficial dorsal horn of the spinal cord. Nonetheless, the functional significance of nAChRs in the deep dorsal horn of adult animals remains unclear. Using whole-cell patch-clamp recordings from lamina V neurons in the adult rat spinal cord, we investigated whether the activation of nAChRs could modulate the inhibitory synaptic transmission in the deep dorsal horn. In the presence of CNQX and APV to block excitatory glutamatergic synaptic transmission, bath applications of nicotine (100  $\mu$ M) significantly increased the frequency of spontaneous inhibitory postsynaptic currents (sIPSCs) in almost all neurons tested. The effect of nicotine was mimicked by N-methyl-4-(3-pyridinyl)-3-butene-1-amine (RJR-2403, 100  $\mu$ M), an  $\alpha$ 4 $\beta$ 2-nAChR agonist, and was also mimicked by choline (10 mM), an  $\alpha$ 7-nAChR agonist. The effect of nicotine was completely blocked by the nAChR antagonist mecamylamine (5  $\mu$ M). In the presence of tetrodotoxin (0.5  $\mu$ M), nicotine (100  $\mu$ M) significantly increased the miniature IPSC frequency. On the other hand, RJR-2403 (100  $\mu$ M) or choline (10 mM) did not affect miniature IPSCs. The application of nicotine (100  $\mu$ M) also evoked a large inward current in all lamina V neurons tested when cells were held at -60 mV. Similarly, RJR-2403 (100  $\mu$ M) induced inward currents in the majority of lamina V neurons examined. On the other hand, choline (10 mM) did not elicit any detectable whole-cell currents. These results suggest that several nAChR subtypes are expressed on the presynaptic terminals, preterminals, and neuronal cell bodies within lamina V and that these nAChRs are involved in the modulation of inhibitory synaptic activity in the deep dorsal horn of the spinal cord.

### Background

Neuronal nAChRs are a larger family of ligand-gated ion channels widely expressed in both the central and the

peripheral nervous system. At least 12 different subunits of nAChRs, including  $\alpha$ 2- $\alpha$ 10,  $\beta$ 2- $\beta$ 4, have been identified so far and these subunits form many different sub-

types of nAChRs with pentameric structures consisting of homomers or heteromers [1]. Homomeric nAChRs are made up of  $\alpha 7$ ,  $\alpha 8$  or  $\alpha 9$  subunits, while heteromeric nAChRs comprise various combinations of  $\alpha 2$ – $\alpha 6$  with  $\beta 2$ – $\beta 4$  subunits,  $\alpha 9$  with  $\alpha 10$  subunits [2,3]. These subtypes of nAChRs have different pharmacological and biophysical properties [1]. It has been shown that nAChRs are involved in a variety of physiological functions including learning, reinforcement, development, aging and nociception [4].

Although Davis et al. (1932) first reported that nicotine has analgesic effects [5], high dosages of nicotine were required to produce antinociception and its effect was relatively modest with a short duration [6-8]. Epibatidine, a potent nAChR agonist isolated from the skin of an Ecuadorian frog, was about 100-fold more potent than morphine in rodents [9-12]. Unfortunately, the dosage of epibatidine to produce antinociception was near that to cause seizure, death, and other side effects [12]. The intolerable toxic effects of epibatidine were due to its actions on a broad range of nAChR subtypes. Therefore, the key to the development of safe and effective nicotinic agonists as analgesics is to first understand which nAChR subtypes are involved in modulating nociceptive transmission.

The spinal dorsal horn is the first site in the central nervous system where somatosensory information is processed and integrated. Multiple subtypes of nAChRs are expressed in the spinal dorsal horn and these receptors have been indicated to modulate sensory inputs from the periphery. Genzen and McGehee (2003) have demonstrated that the activation of  $\alpha 7$  nAChRs located at the central terminals of primary afferents enhances the glutamatergic excitatory transmission in the spinal dorsal horn [13]. Several subtypes of nAChRs have been shown to exert tonic or phasic control on the descending inhibitory serotonergic transmission [14]. Multiple subtypes of nAChRs are found to be expressed on both inhibitory and excitatory interneurons in the spinal dorsal horn [15]. The activation of presynaptic nAChRs facilitates GABAergic and glycinergic inhibitory synaptic transmission in the superficial dorsal horn [16-19]. Although the roles of presynaptic nAChRs were extensively studied in the superficial dorsal horn, it is unclear whether nAChRs also mediate sensory modulation in the deep dorsal horn of the spinal cord in adult animals. A variety of sensory inputs, including nociceptive and non-nociceptive inputs, are transmitted into deep dorsal horn [20]. Deep dorsal horn neurons, especially those in the lamina V region, can generate long-lasting afterdischarges in response to nociceptive inputs and this hyperactivity has important implications in pathological pain states [21]. Inhibitory modulation in this region is critical in preventing the central hyperactivity and hyperalgesia. The aim of this study


was to evaluate the effects of nAChR activation on the inhibitory synaptic transmission in deep dorsal horn neurons.

## Results

### Effects of nicotine and nAChR agonists on spontaneous IPSCs in the lamina V neurons

Whole-cell patch-clamp recordings were performed from lamina V neurons of spinal cord slices prepared from adult rats. Stable recordings could be obtained from slices maintained *in vitro* for more than 12 hours. Glutamatergic excitatory postsynaptic transmission was blocked by CNQX (20  $\mu$ M) and APV (50  $\mu$ M). All lamina V neurons tested exhibited spontaneous inhibitory postsynaptic currents (sIPSCs) when cells were held at -10 mV. In the presence of bicuculline (20  $\mu$ M) and strychnine (2  $\mu$ M), sIPSCs were completely abolished in all lamina V neurons tested ( $n = 3$ ; data not shown), indicating these sIPSCs were mediated by GABA and/or glycine receptors. Perfusion of nicotine (100  $\mu$ M) for 1 min resulted in a rapid and significant increase in sIPSC frequency in all neurons tested (Fig. 1A–C). The average sIPSC frequency in controls was  $2.1 \pm 0.6$  Hz (0.4 – 5.1 Hz,  $n = 9$ ) and the frequency increased to  $15.8 \pm 2.3$  Hz (4.2 – 28.1 Hz,  $n = 9$ ,  $P < 0.05$ ) following the application of 100  $\mu$ M nicotine (Fig. 1C); the sIPSC frequency increased to  $1330 \pm 310\%$  of the control ( $n = 9$ ,  $P < 0.05$ ). The nicotine-induced increase in sIPSC frequency was completely blocked in the presence of nAChR antagonist mecamylamine (5  $\mu$ M) administered 5 min prior to the application of nicotine ( $n = 3$ ; Fig. 1D, E). After the washout of mecamylamine, a second application of nicotine (100  $\mu$ M) increased the sIPSC frequency in all neurons tested (Fig. 1D, E).

We tested RJR-2403, a selective  $\alpha 4\beta 2$  nAChR agonist, to see if it also increased sIPSC frequency. Similar to nicotine, application of 100  $\mu$ M RJR-2403 for 1 min markedly increased sIPSC frequency in 13 out of 14 neurons recorded (Fig. 2Aa–c). The average sIPSC frequency in the control and following the application of RJR-2403 was  $5.8 \pm 1.0$  Hz (0.5 – 11.6 Hz,  $n = 14$ ) and  $15.7 \pm 1.9$  Hz (3.7 – 25.2 Hz,  $n = 14$ ), respectively (Fig. 2Ac). The sIPSC frequency following the applications of RJR-2403 significantly increased to  $573 \pm 189\%$  of the control ( $n = 14$ ,  $P < 0.05$ ; Fig. 2C). While RJR-2403 alone produced a significant increase in sIPSC frequency, the effects of RJR-2403 was completely blocked in the presence of dihydro-beta-erythroidine (Dh $\beta$ E, 1  $\mu$ M), an  $\alpha 4\beta 2$  nAChR antagonist ( $97 \pm 2\%$  of control,  $n = 3$ ). Perfusion of choline (10 mM), a selective  $\alpha 7$  nAChR agonist, for 1 min also increased the sIPSC frequency in 11 out of the 13 neurons examined (Fig. 2Ba–c). The average sIPSC frequency in the control and following the application of choline was  $7.4 \pm 1.2$  Hz (2.9 – 16.9 Hz,  $n = 13$ ) and  $15.2 \pm 2.3$  Hz (5.2 – 27.6 Hz,  $n = 13$ ), respectively (Fig. 2Bc). The sIPSC frequency fol-



**Figure 1**  
**Effects of nicotine on spontaneous IPSCs in lamina V neurons.** **A**, A continuous recording of sIPSCs in the control and following the application of nicotine (Nic, 100  $\mu$ M). **B**, A histogram shows the time course of the changes in sIPSC frequency following the application of nicotine; time bin is 10 s. **C**, The graph shows the individual result from 9 lamina V neurons. **D**, Effects of mecamylamine (Mec) on nicotine-induced increase in sIPSC frequency. The consecutive traces on the left are sIPSCs in the control (upper panel) and following the application of nicotine (lower panel) in the presence of mecamylamine (5  $\mu$ M). The consecutive traces on the right are sIPSCs in the control (upper panel) and following the application of nicotine (lower panel) after the washout of mecamylamine. Note that the bath application of nicotine did not affect the sIPSCs in the presence of mecamylamine, but it markedly increased sIPSC frequency after the washout of mecamylamine. **E**, Two histograms show time courses of changes in sIPSC frequency following the application of nicotine in the presence of mecamylamine (left) and after the washout of mecamylamine (right); time bin is 10 s.


lowing the applications of choline significantly increased to  $221 \pm 22\%$  of control ( $n = 13$ ,  $P < 0.05$ ; Fig. 2C). While choline alone produced a significant increase in sIPSC frequency, choline did not produce any significant increase in sIPSC frequency in the presence of methyllycaconitine (MLA, 50 nM), an  $\alpha 7$  nAChR antagonist ( $98 \pm 2\%$  of control,  $n = 4$ ).

#### Effects of nicotine and nAChR agonists on mIPSCs in lamina V neurons


We examined the effects of nicotine or nAChR agonists on mIPSC frequency in the presence of tetrodotoxin (TTX, 0.5  $\mu$ M) to determine whether nAChRs might be localized at the presynaptic terminals of GABAergic and/or glycinergic inhibitory interneurons. Application of 0.5  $\mu$ M TTX itself blocked the action potential-driven synaptic transmission and decreased the amplitude of IPSCs from  $149.9 \pm 125.6$  pA to  $30.0 \pm 18.4$  pA ( $n = 4$ ). Under this condition, bath application of nicotine (100  $\mu$ M) largely increased mIPSC frequency in all neurons recorded, but there was no effect on mIPSC amplitude ( $n = 10$ ; Fig. 3A). The average mIPSC frequency in the control and following the applications of nicotine was  $1.7 \pm 0.4$  Hz (0.4 – 5.3 Hz,  $n = 10$ ) and  $15.0 \pm 2.3$  Hz (1.4 – 25.2 Hz,  $n = 10$ ), respectively. The mIPSC frequency following the applications of nicotine significantly increased to  $1043 \pm 153\%$  of control ( $n = 10$ ,  $P < 0.05$ , Fig. 3D). On the other hand, perfusion of 100  $\mu$ M RJR-2403 ( $n = 6$ ) or 10 mM choline ( $n = 7$ ) did not affect mIPSC frequency and amplitude (Fig. 3B, 3C). The average mIPSC frequency following the application of RJR-2403 and choline was  $95 \pm 3\%$  of control ( $n = 6$ ) and  $98 \pm 2\%$  of control ( $n = 7$ ), respectively (Fig. 3D).

#### Whole-cell currents directly evoked by nicotine or nAChR agonists in lamina V neurons

We determined whether nicotine, RJR-2403 and choline could evoke whole-cell currents in lamina V neurons. In this set of experiments, cells were held at -60 mV and recordings were conducted in the presence of 20  $\mu$ M CNQX, 50  $\mu$ M APV, 20  $\mu$ M bicuculline and 10  $\mu$ M PMBA (3-[2'-Phosphonomethyl][1,1'-biphenyl]-3-yl)alanine). Under this condition, both excitatory and inhibitory postsynaptic currents were completely disappeared. The bath application of nicotine (100  $\mu$ M) for 1 min evoked an inward current in all neurons tested (Fig. 4A). The average peak amplitude of the inward currents evoked by nicotine was  $95 \pm 19$  pA ( $n = 8$ ; Fig. 4D). The bath application of RJR-2403 (100  $\mu$ M) for 1 min also evoked large inward currents in 6 out of 8 neurons examined (Fig. 4B). The average peak amplitude of the inward currents induced by RJR-2403 was  $119 \pm 42$  pA ( $n = 6$ ; Fig. 4D). In contrast to nicotine and RJR-2403, choline (10 mM) did not elicit any detectable currents ( $n = 6$ ; Fig. 4C).



**Figure 2**  
**Effects of nicotinic receptor agonists on sIPSCs in lamina V neurons.** **A, (a)** A continuous recording of sIPSCs in the control and following the application of the selective  $\alpha 4\beta 2$  nAChR agonist RJR-2403 (100  $\mu$ M). **(b)** A histogram shows the time course of changes in sIPSC frequency following the application of RJR-2403; time bin is 10 s. **(c)** The graph shows the individual result from 14 lamina V neurons. **B,** The experiment was similar to that shown in **(A)** except that the selective  $\alpha 7$  nAChR agonist choline (10 mM) was tested. Similar results were obtained in 11 out of 13 neurons. **C,** A histogram shows relative sIPSC frequency following the application of nicotine ( $n = 9$ ), RJR-2403 ( $n = 14$ ), or choline ( $n = 13$ ). sIPSC frequency before the applications of testing drugs is used as control and is scaled at 100%. Data represent Mean  $\pm$  SEM; \* $P < 0.05$ .

**Figure 3**


**Effects of nicotinic receptor agonists on mIPSCs in lamina V neurons.** **A**, The consecutive traces of mIPSCs are in the control (left) and following the application of nicotine (100  $\mu$ M, right). Two histograms on the bottom panel show the time course of changes in mIPSC frequency (left) and amplitude (right) during the application of nicotine; time bin is 10 s. **B**, **C**, Experiments were similar to that shown in (**A**) except that 100  $\mu$ M RJR-2403 or 10 mM choline was tested. **D**, A histogram shows relative mIPSC frequency following the applications of nicotine ( $n = 11$ ), RJR-2403 ( $n = 6$ ), or choline ( $n = 7$ ). mIPSC frequency before the applications of testing drugs is used as control and is scaled at 100%. Data represent Mean  $\pm$  SEM; \* $P < 0.05$ ; n.s., not significant.

## Discussion

The present study demonstrated in lamina V neurons of the adult rat spinal cord that nicotine increased sIPSC frequency when glutamatergic excitatory transmission was blocked in the presence of CNQX and APV and that nicotine also increased mIPSC frequency when action potential-driven synaptic transmission was not permitted in the presence of TTX. Interestingly, however, neither the  $\alpha 4\beta 2$  nor  $\alpha 7$  nAChR agonists increased mIPSC frequency although both of them increased sIPSC frequency in lamina V neurons. Together with the findings of our previous study conducted on superficial laminae of the spinal cord of adult rats [18], we have provided electrophysiological evidence showing that inhibitory synaptic activity in both superficial and deep laminae of the spinal cord dorsal horn are modulated by different nAChR subtypes.

Nicotinic receptors are abundant in different CNS regions, where they are shown to regulate the release of various


neurotransmitters, including serotonin, norepinephrine, glutamate, GABA and glycine [22-24]. In the present study, the activation of nAChRs enhanced the GABA and/or glycine release onto lamina V neurons. A similar enhancement of the inhibitory synaptic transmission by the activation of nAChRs has been reported in the superficial layers of the spinal dorsal horn [16,18,19]. In neonatal rats,  $\alpha 4\beta 2$  nAChR subtype has been suggested to be expressed at presynaptic terminals and these receptors mediate significantly increases in the glycinergic [16] and GABAergic inhibitory synaptic transmission in the superficial lamina of the spinal cord dorsal horn [19]. Interestingly, the expression of nAChR subunits in the spinal cord changes during development [25]. Consistent with the development changes of nAChR subunits, our previous study demonstrated that a non- $\alpha 4\beta 2$  and non- $\alpha 7$  subtype nAChR mediated an enhancement of both the GABAergic and glycinergic mIPSC frequency in the superficial laminae of the adult rat spinal dorsal horn [18]. In the deeper



**Figure 4**  
**Whole-cell currents evoked by nicotine and nAChR agonists in lamina V neurons.** **A**, Whole-cell currents evoked by bath application of nicotine (100  $\mu$ M) in a lamina V neuron. **B**, Whole-cell currents evoked by RJR-2403 (100  $\mu$ M) in a different lamina V neuron. **C**, Choline (10 mM) did not evoke any membrane current in a neuron. **D**, The histogram shows the average amplitude of the whole-cell inward currents induced by nicotine ( $n = 8$ ), RJR-2403 ( $n = 6$ ), and choline ( $n = 6$ ).

laminae (lamina III–V) of neonatal rats, a previous study showed that presynaptic  $\alpha 4\beta 2$  nAChRs mediate the facilitation of GABA release [19], a result similar to those shown in superficial laminae of neonatal rats [16,19]. However, the present study revealed that the nAChR-mediated modulation of inhibitory synaptic transmission in the adult stage is more complicated than that in the neonatal stage. We showed that nicotine largely increased both sIPSC frequency and mIPSC frequency in the lamina V neurons in adult rats. On the other hand, RJR-2403, a potent activator of  $\alpha 4\beta 2$  nAChR, and choline, a selective agonist for  $\alpha 7$  nAChR, significantly increased the sIPSC frequency, but did not change mIPSC frequency. It has been demonstrated that nAChRs are expressed at two cellular locations in the central nervous system [26]. One is presynaptic sites or synaptic boutons where nAChR activation modulates transmitter release in a TTX-insensitive manner. The other is preterminals at terminal axon branches where nAChR activation affects transmitter release by depolarizing axonal membranes to fire action potentials. A recent immunohistochemical study revealed the immunoreactivity of nAChRs in lamina V neurons of the spinal dorsal horn at both presynaptic and preterminal sites [27]. At the preterminal sites, nAChR-mediated regulation of transmitter releases is TTX-sensitive and can be blocked in the presence of TTX. The block by TTX of the

RJR-2403- and choline-induced increase in sIPSC frequency in our results suggests that  $\alpha 4\beta 2$  and  $\alpha 7$  nAChRs are not expressed at presynaptic terminals. These receptors are likely to be expressed at the preterminals or other parts of GABAergic and/or glycinergic neurons whose axons innervate lamina V neurons. Because nicotine could still increase IPSC frequency in the presence of TTX, it suggests that a non- $\alpha 4\beta 2$ , non- $\alpha 7$  subtype of nAChR is located at the presynaptic terminals of GABAergic and/or glycinergic neurons that innervate lamina V neurons (Fig. 5). The



**Figure 5**  
**Schematic diagram of nAChR-mediated modulation of sensory synaptic transmission in the dorsal horn of adult rats.** In adult rats,  $\alpha 4\beta 2$  nAChRs are expressed on the soma of inhibitory interneurons located in the lamina V region. These nAChRs may be also expressed on preterminals but not at presynaptic sites of the lamina V inhibitory neurons. Lamina V neurons are also synapsed by inhibitory neurons expressing  $\alpha 7$  nAChRs at their preterminals and/or on their somas and these inhibitory neurons are likely to be located in other lamina regions. The modulation of inhibitory activity in lamina V by both  $\alpha 4\beta 2$  nAChRs and  $\alpha 7$  nAChRs depends on membrane depolarization and action potentials. There is a non- $\alpha 4\beta 2$ /non- $\alpha 7$  subtype of nAChRs that are expressed at the presynaptic terminals of inhibitory neurons in lamina V region. The modulation of inhibitory transmission in lamina V by non- $\alpha 4\beta 2$ /non- $\alpha 7$  subtype of nAChRs is independent of membrane depolarization and action potentials. The distribution of nAChR subtypes in the spinal cord lamina II region of adult rats [18] is also presented in this diagram for a comparison.

presence of a non- $\alpha 4\beta 2$ , non- $\alpha 7$  subtype of nAChR in spinal cord dorsal horns are supported by previous studies using  $\alpha 4\beta 2$  nAChR knock-out mouse [28], *in situ* hybridization [29] and the combination of patch-clamp recordings with single-cell RT-PCR [15]. All these previous studies pointed to the potential presence of other functional nAChR subtypes in addition to  $\alpha 4\beta 2$  and  $\alpha 7$  receptors. However, the subunit compositions of the non- $\alpha 4\beta 2$  and non- $\alpha 7$  subtype of nAChRs remain to be identified.

We have shown that both nicotine and the  $\alpha 4\beta 2$  nAChR agonist RJR-2403 directly evoked whole-cell inward currents in the majority of lamina V neurons. On the other hand, the  $\alpha 7$  nAChR agonist choline did not evoke any detectable inward currents in lamina V neurons. Since strychnine has been noted to also be an effective antagonist at  $\alpha 7$  nAChRs [30,31], the effect of choline was examined in the presence of PMBA, a glycine receptor antagonist that has no effect on  $\alpha 7$  nAChRs [32,33]. The lack of choline-evoked whole-cell currents raise a possibility that  $\alpha 7$  nAChRs are not expressed on lamina V neurons. If this is the case, then choline-induced increases of sIPSC frequency might be mediated by  $\alpha 7$  nAChRs that are expressed on the inhibitory interneurons whose cell bodies are located in other lamina regions in the spinal cord. In contrast to  $\alpha 7$  nAChRs, our results suggest that  $\alpha 4\beta 2$  nAChR expressing inhibitory interneurons are located in lamina V. Consistently, several reports have indicated the presence of  $\alpha 4\beta 2$  nAChR in the deep dorsal horn and  $\alpha 7$  nAChRs in other lamina of the spinal cord [15,33-35]. Bradia et al. (2002) has reported that  $\alpha$ -bungarotoxin-sensitive  $\alpha 7$  nAChRs are located in the parasympathetic preganglionic neurons surrounding the central canal of the spinal cord (lamina X) [33]. A low level of  $\alpha 7$  transcripts were also detected by *in situ* hybridization in the area around the central canal [35]. Moreover, the single-cell RT-PCR study revealed a more widespread expression of  $\alpha 7$  nAChR subunits in mouse spinal dorsal horn neurons [15]. These findings support the idea that  $\alpha 7$  nAChR-expressing inhibitory interneurons innervate lamina V neurons from other lamina regions in the spinal cord.

The role of nAChRs in modulating pain transmission has been reported by a number of studies. Using  $\alpha 4\beta 2$  knock-out mice, Marubio et al. (1999) showed a reduced antinociceptive effect in a behavior study [28]. In a neuropathic mouse model, epibatidine, a potent agonist of nAChRs showed strong analgesic effects. However, the effects of epibatidine were not completely prevented by the  $\alpha 4\beta 2$  nAChR antagonist dihydro- $\beta$ -erythroidine [36]. These studies suggested that in addition to  $\alpha 4\beta 2$  nAChR, other nAChRs were involved in nAChR-mediated analgesic effects. Consistent with this idea, the intrathecal injection of choline, an  $\alpha 7$  nAChR agonist, has been reported to

have an antinociceptive effect [37]. Several mechanisms have been proposed to contribute to nAChR-mediated analgesic effects, including the desensitization of nAChRs on nociceptive primary afferent fibers, the increase of noradrenaline and serotonin release within the spinal cord, the activation of the descending inhibitory pathways [14,38], and the increases of GABA and glycine release from inhibitory interneurons in the superficial spinal cord dorsal horn [39]. Our study suggest that  $\alpha 4\beta 2$ ,  $\alpha 7$  nAChRs, and another undefined subtype of nAChRs are involved in regulating GABA and/or glycine release in the deep lamina of the spinal cord dorsal horn in adult animals.

In this study, the application of nicotine or nicotinic agonists significantly facilitated GABAergic and/or glycinergic inhibitory synaptic transmission in the deep dorsal horn of the spinal cord. This raises a possibility that acetylcholine released endogenously may modulate inhibitory synaptic transmission in a similar fashion. Recently, Rashid et al. (2006) suggested that endogenous acetylcholine tonically stimulated the GABA and glycine release via  $\alpha 4\beta 2$  subtype of nAChRs in the superficial dorsal horn in mice [40]. Acetylcholine may be released from the interneurons in the dorsal horn since the cell bodies of cholinergic interneurons have been found in lamina III-V [41]. It appears that in the deep dorsal horn there are no descending cholinergic systems in the rat [20,42]. Thus, cholinergic interneurons in the dorsal horn [43,44] may play an important role in modulating inhibitory synaptic transmission.

In the dorsal horn, GABAergic and glycinergic inhibitory synapses undergo developmental changes [45-47]. In the present study, we did not separate inhibitory activity between those of GABAergic synapses and those of glycinergic synapses. It would be interesting to further study whether nAChR subtype expression on GABAergic and glycinergic neurons is different in the spinal cord dorsal horn.

## Conclusion

We have demonstrated that several nAChR subtypes are expressed on the presynaptic terminals, preterminals, and neuronal cell bodies within lamina V and that they are involved in the facilitation of inhibitory synaptic transmission. Therefore, the activation of nAChRs in the deep dorsal horn of the spinal cord may be capable of inhibiting nociceptive signaling in physiological and pathological pain sensations.

## Methods

All the experimental procedures involving the use of animals were approved by the Ethics Committee on Animal Experiments, Wakayama Medical University, and were in

accordance with the UK Animals (Scientific Procedures) Act 1986 and all associated guidelines.

### Spinal cord slice preparation

The method used to prepare adult rat spinal cord slices has been described previously [48]. In brief, male adult Sprague-Dawley rats (6–8 weeks of age, 200–300 g) were deeply anaesthetized with isoflurane through a nose cone inhalation, and then lumbosacral laminectomy was performed. The lumbosacral spinal cord (L1-S3) was removed and placed in pre-oxygenated Krebs solution at 1–3°C. Immediately after the removal of the spinal cord, the rats were killed by exsanguination. The pia-arachnoid membrane was removed after cutting all the ventral and dorsal roots near the root entry zone. The spinal cord was mounted on a vibratome and then a 600 µm-thick transverse slice was cut. The slice was placed on nylon mesh in the recording chamber, which had a volume of 0.5 ml, and then was perfused at a rate of 10–15 ml/min with Krebs solution saturated with 95% O<sub>2</sub> and 5% CO<sub>2</sub>, and maintained at room temperature. A platinum grid was placed on the top of the slice to prevent slice movement. The Krebs solution contained (in mM) 117 NaCl, 3.6 KCl, 2.5 CaCl<sub>2</sub>, 1.2 MgCl<sub>2</sub>, 1.2 NaH<sub>2</sub>PO<sub>4</sub>, 25 NaHCO<sub>3</sub> and 11 glucose.

### Patch-clamp recordings from lamina V neurons

Blind whole-cell patch-clamp recordings were made from lamina V neurons with patch-pipette electrodes having a resistance of 5–10 MΩ [48]. The patch-pipette solution was composed of (in mM) 110 Cs<sub>2</sub>SO<sub>4</sub>, 5 Tetraethylammonium (TEA), 0.5 CaCl<sub>2</sub>, 2 MgCl<sub>2</sub>, 5 EGTA, 5 HEPES, 5 ATP-Mg, pH 7.2. Signals were acquired with a patch-clamp amplifier (Axopatch 200B; Axon Instruments, Foster City, CA, USA). The data were digitized with an A/D converter (Digidata 1200, Axon Instruments) and stored and analyzed with a personal computer using the pCLAMP data acquisition program (Version 8.2, Axon Instruments). Lamina V neurons were viable for up to 24 h in slices perfused with a pre-oxygenated Krebs solution. All the recordings described in this study were made within 12 h. Whole-cell patch-clamp recordings were stable for up to 4 h. All of the neurons had membrane potentials more negative than -50 mV. Unless otherwise noted, all the recordings in this study were performed in the presence of CNQX (20 µM) and APV (50 µM).

### Drug Applications

Drugs were dissolved in Krebs solution and then were applied by perfusion via a three-way stopcock without any change in the perfusion rate or the temperature. The time necessary for the solution to flow from the stopcock to the surface of the spinal cord slice was approximately 20 s. The drugs used in this study were nicotine (Sigma-Aldrich, St. Louis, MO, USA), RJR2403 (Tocris, Ballwin, MO,

USA), choline (Sigma-Aldrich), mecamylamine (Sigma-Aldrich), 3-[2'-Phosphonomethyl[1,1'-biphenyl]-3-yl]alanine (PMBA, Sigma RBI), bicuculline (Sigma-Aldrich), strychnine (Sigma-Aldrich), 6-cyano-7-nitroquinoxaline-2,3-dion (CNQX, Tocris), D(-)-2-Amino-5-phosphonopentanoic acid (D-APV, Tocris), and tetrodotoxin (TTX, Tocris).

### Statistical analysis

All numerical data were expressed as the mean ± S.E.M. Statistical significance was determined as P < 0.05 using paired Student's t-test. For electrophysiological data, *n* refers to the number of neurons recorded.

### Abbreviations

nACh, nicotinic acetylcholine receptor;

IPSC, inhibitory postsynaptic current;

RJR2-403, N-methyl-4-(3-pyridinyl)-3-butene-1-amine;

TTX, tetrodotoxin;

GABA, gamma-aminobutyric acid;

TEA, tetraethylammonium;

EGTA, ethyleneglycol bis(2-aminoethylether)tetraacetic acid;

HEPES, N-(2-Hydroxyethyl)piperazine-N'-(2-ethanesulfonic acid);

ATP-Mg, adenosine triphosphate-magnesium;

CNQX, 6-cyano-7-nitroquinoxaline-2,3-dion;

PMBA, 3-[2'-Phosphonomethyl[1,1'-biphenyl]-3-yl]alanine;

D-APV, D(-)-2-Amino-5-phosphonopentanoic acid;

### Competing interests

The author(s) declare that they have no competing interests.

### Acknowledgements

This work was supported by The General Insurance Association of Japan, The Japanese Health Sciences Foundation, and Grants-in-Aid for Scientific Research from the Ministry of Education, Science, Sports and Culture of Japan to T.N.

### References

1. Changeux JP, Edelstein SJ: **Allosteric receptors after 30 years.** *Neuron* 1998, **21**:959-980.
2. McGehee DS: **Molecular diversity of neuronal nicotinic acetylcholine receptors.** *Ann N Y Acad Sci* 1999, **868**:565-577.



3. Dani JA, Ji D, Zhou FM: **Synaptic plasticity and nicotine addiction.** *Neuron* 2001, **31**:349-352.
4. Cordero-Erausquin M, Marubio LM, Klink R, Changeux JP: **Nicotinic receptor function: new perspectives from knockout mice.** *Trends Pharmacol Sci* 2000, **21**:211-217.
5. Davis L, Pollock LJ, Stone TT: **Visceral pain.** *Surg Gynecol Obstet* 1932, **55**:418-427.
6. Sahley TL, Berntson GG: **Antinociceptive effects of central and systemic administrations of nicotine in the rat.** *Psychopharmacology* 1979, **65**:279-283.
7. Tripathi HL, Martin BR, Aceto MD: **Nicotine-induced antinociception in rats and mice: correlation with nicotine brain levels.** *J Pharmacol Exp Ther* 1982, **221**:91-96.
8. Aceto MD, Aways H, Martin BR, May EL: **Antinociceptive action of nicotine and its methiodide derivatives in mice and rats.** *Br J Pharmacol* 1983, **79**:869-876.
9. Spande TF, Garraffo HM, Yeh HJ, QL, Pannell LK, Daly JW: **A new class of alkaloids from a dendrobatid poison frog: a structure for alkaloid 251F.** *J Nat Prod* 1992, **55**:707-722.
10. Qian C, Li T, Shen TY, Libertine-Garahan L, Eckman J, Biftu T, Ip S: **Epibatidine is a nicotinic analgesic.** *Eur J Pharmacol* 1993, **250**:R13-14.
11. Badio B, Daly JW: **Epibatidine, a potent analgesic and nicotinic agonist.** *Mol Pharmacol* 1994, **45**:563-569.
12. Sullivan JP, Decker MW, Brioni JD, Donnelly-Roberts D, Anderson DJ, Bannon AW, Kang CH, Adams P, Piattoni-Kaplan M, Buckley MJ, et al.: **(+/-)-Epibatidine elicits a diversity of in vitro and in vivo effects mediated by nicotinic acetylcholine receptors.** *J Pharmacol Exp Ther* 1994, **271**:624-631.
13. Genzen JR, McGehee DS: **Short- and long-term enhancement of excitatory transmission in the spinal cord dorsal horn by nicotinic acetylcholine receptors.** *Proc Natl Acad Sci USA* 2003, **100**:6807-6812.
14. Cordero-Erausquin M, Changeux JP: **Tonic nicotinic modulation of serotonergic transmission in the spinal cord.** *Proc Natl Acad Sci USA* 2001, **27**:2803-2807.
15. Cordero-Erausquin M, Pons S, Faure P, Changeux JP: **Nicotine differentially activates inhibitory and excitatory neurons in the dorsal spinal cord.** *Pain* 2004, **109**:308-318.
16. Kiyosawa A, Katsurabayashi S, Akaiki N, Pang ZP: **Nicotine facilitates glycine release in the rat spinal dorsal horn.** *J Physiol* 2001, **536**:101-110.
17. Fucile S, Lax P, Eusebi F: **Nicotine modulates the spontaneous synaptic activity in cultured embryonic rat spinal cord interneurons.** *J Neurosci Res* 2002, **67**:329-336.
18. Takeda D, Nakatsuka T, Papke R, Gu JG: **Modulation of inhibitory synaptic activity by a non- $\alpha 4\beta 2$ , non- $\alpha 7$  subtype of nicotinic receptors in the substantia gelatinosa of adult rat spinal cord.** *Pain* 2003, **101**:13-23.
19. Genzen JR, McGehee DS: **Nicotinic modulation of GABAergic synaptic transmission in the spinal cord dorsal horn.** *Brain Res* 2005, **1031**:229-237.
20. Willis WD, Coggeshall RE: *Sensory Mechanisms of the Spinal Cord* 2nd edition. New York: Plenum; 1991.
21. Woolf CJ, King AE: **Physiology and morphology of multireceptive neurons with C-afferent fiber inputs in the deep dorsal horn of the rat lumbar spinal cord.** *J Neurophysiol* 1987, **58**:460-479.
22. Singer S, Rossi S, Verzosa S, Hashim A, Lonow R, Cooper T, Sershen H, Lajtha A: **Nicotine-induced changes in neurotransmitter levels in brain areas associated with cognitive function.** *Neurochem Res* 2004, **29**(9):1779-92.
23. McGehee DS, Heath MJ, Gelber S, Devay P, Role LW: **Nicotine enhancement of fast excitatory synaptic transmission in CNS by presynaptic receptors.** *Science* 1995, **269**:1692-1696.
24. Wang J, Wang X, Irnaten M, Venkatesan P, Evans C, Baxi S, Mendelowitz D: **Endogenous acetylcholine and nicotine activation enhances GABAergic and glycinergic inputs to cardiac vagal neurons.** *J Neurophysiol* 2003, **89**(5):2473-81.
25. Keiger CJ, Prevette D, Conroy WG, Oppenheim RW: **Developmental expression of nicotinic receptors in the chick and human spinal cord.** *J Comp Neurol* 2003, **455**:86-99.
26. Lena C, Changeux JP, Mulle C: **Evidence for "preterminal" nicotinic receptors on GABAergic axons in the rat interpeduncular nucleus.** *J Neurosci* 1993, **13**:2680-2688.
27. Vincler M, Eisenach JC: **Plasticity of spinal nicotinic acetylcholine receptors following spinal nerve ligation.** *Neurosci Res* 2004, **48**(2):139-45.
28. Marubio LM, del Mar Arroyo-Jimenez M, Cordero-Erausquin M, Lena C, Le Novere N, de Kerchove d'Exaerde A, Huchet M, Damaj MI, Changeux JP: **Reduced antinociception in mice lacking neuronal nicotinic receptor subunits.** *Nature* 1999, **398**:805-810.
29. Wada E, Wada K, Boulter J, Deneris E, Heinemann S, Patrick J, Swanson LW: **Distribution of  $\alpha 2$ ,  $\alpha 3$ ,  $\alpha 4$ , and  $\beta 2$  neuronal nicotinic receptor subunit mRNAs in the central nervous system: a hybridization histochemical study in the rat.** *J Comp Neurol* 1989, **284**:314-335.
30. Matsubayashi H, Alkondon M, Pereira EF, Swanson KL, Albuquerque EX: **Strychnine: a potent competitive antagonist of  $\alpha$ -bungarotoxin-sensitive nicotinic acetylcholine receptors in rat hippocampal neurons.** *J Pharmacol Exp Ther* 1998, **284**:904-913.
31. Garcia-Colunga J, Mileti R: **Modulation of nicotinic acetylcholine receptors by strychnine.** *Proc Natl Acad Sci USA* 1999, **30**:4113-4118.
32. Saitoh T, Ishida M, Maruyama M, Shinozaki H: **A novel antagonist, phenylbenzene omega-phosphono-alpha-amino acid, for strychnine-sensitive glycine receptors in the rat spinal cord.** *Br J Pharmacol* 1994, **113**:165-170.
33. Bradaia A, Trouslard J: **Fast synaptic transmission mediated by  $\alpha$ -bungarotoxin-sensitive nicotinic acetylcholine receptors in lamina X neurones of neonatal rat spinal cord.** *J Physiol* 2002, **544**:727-739.
34. Urban L, Willetts J, Murase K, Randic M: **Cholinergic effects on spinal dorsal horn neurons in vitro: an intracellular study.** *Brain Res* 1989, **500**:12-20.
35. Séguéla P, Wadiche J, Dineley-Miller K, Dani JA, Patrick JW: **Molecular cloning, functional properties, and distribution of rat brain  $\alpha 7$ : a nicotinic cation channel highly permeable to calcium.** *J Neurosci* 1993, **13**:596-604.
36. Rashid MH, Ueda H: **Neuropathy-specific analgesic action of intrathecal nicotinic agonists and its spinal GABA-mediated mechanism.** *Brain Res* 2002, **953**:53-62.
37. Damaj MI, Meyer EM, Martin BR: **The antinociceptive effects of  $\alpha 7$  nicotinic agonists in an acute pain model.** *Neuropharmacol* 2000, **39**:2785-2791.
38. Li X, Eisenach JC: **Nicotinic acetylcholine receptor regulation of spinal norepinephrine release.** *Anesthesiology* 2002, **96**:1450-1456.
39. Todd AJ, Spike RC: **The localization of classical transmitters and neuropeptides within neurons in laminae I-III of the mammalian spinal dorsal horn.** *Prog Neurobiol* 1993, **41**:609-645.
40. Rashid MH, Furue H, Yoshimura M, Ueda H: **Tonic inhibitory role of  $\alpha 4\beta 2$  subtype of nicotinic acetylcholine receptors on nociceptive transmission in the spinal cord in mice.** *Pain* 2006, **125**(1-2):125-35.
41. Olave MJ, Puri N, Kerr R, Maxwell DJ: **Myelinated and unmyelinated primary afferent axons form contacts with cholinergic interneurons in the spinal dorsal horn.** *Exp Brain Res* 2002, **145**(4):448-56.
42. Bowker RM, Westlund KN, Sullivan MC, Wilber JF, Coulter JD: **Descending serotonergic, peptidergic and cholinergic pathways from the raphe nuclei: a multiple transmitter complex.** *Brain Res* 1983, **288**:33-48.
43. Barber RP, Phelps PE, Houser CR, Crawford GD, Salvaterra PM, Vaughn JE: **The morphology and distribution of neurons containing choline acetyltransferase in the adult rat spinal cord: an immunocytochemical study.** *J Comp Neurol* 1984, **229**:329-346.
44. Todd AJ: **Immunohistochemical evidence that acetylcholine and glycine exist in different populations of GABAergic neurons in lamina III of rat spinal dorsal horn.** *Neuroscience* 1991, **44**:741-746.
45. Baccei ML, Fitzgerald M: **Development of GABAergic and glycinergic transmission in the neonatal rat dorsal horn.** *J Neurosci* 2004, **24**(20):4749-57.
46. Allain AE, Bairo A, Meyrand P, Branchereau P: **Expression of the glycinergic system during the course of embryonic development in the mouse spinal cord and its co-localization with GABA immunoreactivity.** *J Comp Neurol* 2006, **496**(6):832-46.

47. Ataka T, Gu JG: **Relationship between tonic inhibitory currents and phasic inhibitory activity in the spinal cord lamina II region of adult mice.** *Mol Pain* 2006, **2**:2:36.
48. Nakatsuka T, Ataka T, Kumamoto E, Tamaki T, Yoshimura M: **Alteration in synaptic inputs through C-afferent fibers to substantia gelatinosa neurons of the rat spinal dorsal horn during postnatal development.** *Neuroscience* 2000, **99**:549-556.

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

