

RESEARCH

Open Access

Milnacipran inhibits glutamatergic *N*-Methyl-D-Aspartate receptor activity in Spinal Dorsal Horn Neurons

Tatsuro Kohno^{1,4*}, Masafumi Kimura², Mika Sasaki³, Hideaki Obata^{2,4}, Fumimasa Amaya^{3,4} and Shigeru Saito²

Abstract

Background: Antidepressants, which are widely used for treatment of chronic pain, are thought to have antinociceptive effects by blockade of serotonin and noradrenaline reuptake. However, these drugs also interact with various receptors such as excitatory glutamatergic receptors. Thermal hyperalgesia was induced by intrathecal injection of NMDA in rats. Paw withdrawal latency was measured after intrathecal injection of antidepressants. The effects of antidepressants on the NMDA and AMPA-induced responses were examined in lamina II neurons of rat spinal cord slices using the whole-cell patch-clamp technique. The effects of milnacipran followed by application of NMDA on pERK activation were also investigated in the spinal cord.

Results: Intrathecal injection of milnacipran (0.1 μ mol), but not citalopram (0.1 μ mol) and desipramine (0.1 μ mol), followed by intrathecal injection of NMDA (1 μ g) suppressed thermal hyperalgesia. Milnacipran (100 μ M) reduced the amplitude of NMDA (56 ± 3 %, 64 ± 5 % of control)-, but not AMPA (98 ± 5 %, 97 ± 5 % of control)-mediated currents induced by exogenous application and dorsal root stimulation, respectively. Citalopram (100 μ M) and desipramine (30 μ M) had no effect on the amplitude of exogenous NMDA-induced currents. The number of pERK-positive neurons in the group treated with milnacipran (100 μ M), but not citalopram (100 μ M) or desipramine (30 μ M), followed by NMDA (100 μ M) was significantly lower compared with the NMDA-alone group.

Conclusions: The antinociceptive effect of milnacipran may be dependent on the drug's direct modulation of NMDA receptors in the superficial dorsal horn. Furthermore, in addition to inhibiting the reuptake of monoamines, glutamate NMDA receptors are also important for analgesia induced by milnacipran.

Keywords: Antidepressants, *N*-methyl-D-aspartate (NMDA) Receptor, Spinal Analgesia

Background

It is well established that antidepressants have antinociceptive effects; because of this, they are widely used for treatment of chronic pain [1]. In particular, tricyclic antidepressants (TCAs) have long been the mainstay of treatment for neuropathic pain, which is due to lesion or dysfunction of the peripheral or central nervous system. Antidepressants have the unique ability to inhibit the presynaptic reuptake of monoamines, serotonin (5-HT),

and noradrenaline (NA) at the neuronal terminals [2], and this activity can produce antinociceptive effects. Recently, more selective monoamine reuptake inhibitors, such as 5-HT and NA reuptake inhibitors (SNRIs) and selective 5-HT reuptake inhibitors (SSRIs), have been introduced and are clinically used to treat neuropathic pain [1]. However, the underlying mechanisms of these drugs may be more complex than simply the blockade of 5-HT and NA reuptake.

In fact, TCAs could also interact with various receptors including *N*-methyl-D-aspartate (NMDA) receptors to produce nociceptive effects. NMDA glutamate receptors are one of the major receptor channel types mediating rapid excitatory neurotransmission in the central nervous system, and they also play an important role in central sensitization regarding long-term pain [3,4].

* Correspondence: kohno-t@umin.net

¹Department of Anesthesiology, Niigata University Graduate School of Medical and Dental Sciences, 757 Asahimachi, Chuo ku Niigata 951-8510, Japan

⁴Pain Mechanism Research Group, 757 Asahimachi, Chuo ku Niigata 951-8510, Japan

Full list of author information is available at the end of the article

Some previous studies have suggested that there are interactions in nociceptive transmission between TCAs and NMDA receptors in the spinal cord [5]. However, there have been no studies demonstrating that the antinociceptive effects of antidepressants are attributable to the inhibition of NMDA receptors in the spinal cord at the cellular level.

The spinal cord is an important site of action of antidepressant-mediated antinociception [6]. The brainstem-spinal descending 5-HT and NA systems suppress nociceptive signals from primary afferent neurons to the spinal dorsal horn. Concurrently, the superficial dorsal horn preferentially receives nociceptive primary afferent fibers. Thus, the spinal dorsal horn is thought to play an important role in modulating nociceptive transmission from the periphery [7,8] as well as in regulating the antinociceptive effect of antidepressants.

The purpose of the current study was to test the hypothesis that 3 different antidepressants, milnacipran as a SNRI, citalopram as a SSRI, and desipramine as a TCA (a preferential noradrenergic reuptake inhibitor) have a direct antagonistic effect on NMDA and AMPA-mediated responses in the spinal dorsal horn.

Methods

Surgical Preparation

The study was approved by the Animal Care and Use Committee of the Gunma University School of Medicine (Maebashi, Japan). Male rats (250–270 g) were used in all experiments. Animals were housed under a 12-h light–dark cycle with food and water ad libitum. For intrathecal administration, a sterilized 32-gauge polyethylene catheter (ReCathCo, Allison Park, PA) connected to an 8.5-cm Tygon external tubing (Saint-Gobain Performance Plastics, Akron, OH) was inserted under isoflurane anesthesia, as previously described [9]. The catheter was passed caudally 8.0 cm from the cisterna magnum to the lumbar enlargement. The animals were allowed to recover for 1 week before being used experimentally. Only animals without evidence of neurologic dysfunction after catheter insertion were used for all studies.

Testing Procedures

The thermal nociceptive threshold was measured with a device (Plantar Test[®], IITC Inc. Life Science, Woodland Hills, CA) using a method similar to that reported previously [10]. Rats were placed in individual plastic boxes (10 × 20 × 24 cm) on the glass surface of the testing apparatus, which was maintained at 30 °C during all testing, and were allowed to acclimate for 30 min. Paw withdrawal latency (PWL) was determined using an intense light focused on the hind paw, as previously described [10]. Light intensity was adjusted so that baseline latency was between 9 and 11 s in all animals. A cutoff of 20 s

was selected to avoid tissue damage during periods of analgesia, but no animals reached this cutoff point.

The thermal hyperalgesic state was induced by intrathecal injection of 1 µg NMDA. The dose was selected according to a previous study [11]. This experiment is most specific to evaluate the effect of antidepressants on NMDA-mediated responses in the spinal cord. PWL was measured three times in the right or left foot in the middle of the footpad. These three observations were averaged for each animal. PWLs were measured before and after intrathecal injection of milnacipran, citalopram, desipramine, or saline (six animals in each group), and NMDA was injected intrathecally 15 min after the injection (time 0). PWLs were measured at 0, 30, 60, 90, and 120 min after intrathecal injection of NMDA.

Drugs and Their Administration

The agents administered in this study were milnacipran, citalopram, desipramine, and NMDA. Each antidepressant or saline was administered intrathecally 15 min prior to NMDA injection. Drugs were administered intrathecally in a volume of 5 µl, followed by an injection of 10 µl of saline to flush the catheter. All drugs were dissolved in normal saline. The doses of milnacipran were selected according to the previous studies [12,13]. The maximum dose of milnacipran (0.1 µmol) was used for desipramine and citalopram injections because the doses to produce analgesia in these 3 drugs are almost same [14,15].

In Vitro Patch-Clamp Recordings

This portion of the study was approved by the Animal Care and Use Committee at Niigata University Graduate School of Medical and Dental Sciences (Niigata, Japan). Male rats (150–200 g) were anesthetized with urethane (1.5 g/kg, i.p.). A dorsal laminectomy was performed, and the lumbosacral segment of the spinal cord with ventral and dorsal roots attached was removed [16,17]. The rats were then immediately killed by exsanguination, and the spinal cords were placed in pre-oxygenated ice-cold Krebs solution. After the arachnoid membrane was removed, the spinal cord was placed in an agar block and mounted on a metal stage. A transverse slice (500-µm thick) with an attached dorsal root was cut on a DTK-1500 Microslicer (Dosaka, Kyoto, Japan) and placed on a nylon mesh in the recording chamber. The slice was perfused continuously with Krebs solution (10 ml/min) equilibrated with a 95 % O₂ and 5 % CO₂ gas mixture at 36°C. The Krebs solution contained (in mM): NaCl, 117; KCl, 3.6; CaCl₂, 2.5; MgCl₂, 1.2; NaH₂PO₄, 1.2; NaHCO₃, 25; and D-glucose, 11.5. Whole-cell patch-clamp recordings were made from lamina II neurons in voltage-clamp mode using patch pipette electrodes having a resistance of 10 MΩ. The patch pipette solution contained (in mM): Cs-sulfate, 110; CaCl₂, 0.5; MgCl₂, 2; EGTA, 5; HEPES, 5; tetraethylammonium

(TEA), 5; and ATP-Mg salt, 5. Signals were amplified using an Axopatch 200B amplifier (Molecular Devices, Union City, CA) and were low-pass filtered with a 2-kHz cutoff and digitized at 5 kHz. Data were collected and analyzed using pClamp 10.0 software (Molecular Devices). All experiments were performed in voltage-clamp mode at a holding potential of -40 mV for recording exogenous NMDA current and -70 mV for recording exogenous AMPA current. To test the synaptic response, NMDA receptor-mediated excitatory postsynaptic currents (EPSCs) were observed by dorsal root electrical stimulation at $+40$ mV in the presence of an AMPA/kainate receptor antagonist (6-cyano-7-nitroquinoxaline-2, 3-dione, CNQX; 10 μ M), a GABA_A receptor antagonist (bicuculline; 20 μ M), and a glycine receptor antagonist (strychnine; 1 μ M). AMPA receptor-mediated EPSCs were observed by dorsal root stimulation at -70 mV. Evoked EPSCs that displayed a constant latency and lack of failures with high frequency stimulation (20 Hz) were classified as monosynaptic. Drugs were applied by superfusion without alteration of the perfusion rate and temperature. NMDA (100 μ M) and AMPA (10 μ M) were applied to slices for 30 s, and each antidepressant was applied to slices for 3 min. Peak NMDA and AMPA currents were measured before and after each treatment and expressed as (posttreatment/pretreatment) $\times 100$ (as percentages).

Immunohistochemistry

The spinal cord slices (700 μ m) from male rats (150 – 250 g) were perfused with Krebs solution for at least 3 h before application of drugs. Milnacipran (100 μ M) was applied for 10 min before NMDA stimulation (100 μ M for 5 min) and was present during NMDA stimulation. After drug treatment, the slices were fixed in 4% paraformaldehyde for 60 min, replaced with sucrose overnight, cut in a cryostat at a thickness of 16 μ m, and mounted on slide glass. Phosphorylated extracellular signal-regulated kinase 1/2 (pERK1/2) was visualized by indirect immunohistochemistry [18]. Sections were incubated with rabbit anti-pERK1/2 antibody (Cell Signaling Technology, Danvers, MA; $1:1000$) for 2 days at 4 °C. The sections were incubated with biotinylated anti-rabbit secondary antibody (Vector Laboratories, Burlingame, CA; $1:400$) for 4 h at room temperature. Signals were visualized with Vectastain ABC systems (Vector Laboratories) following the manufacturer's instruction.

Signals were analyzed under a microscope-digital camera system (Nikon, Tokyo, Japan). Experimenters who were unaware of the experimental protocol counted cells in a blinded manner. At least five nonadjacent sections were randomly selected for cell count. The number of pERK1/2-positive neurons in the superficial dorsal horn from each of five sections was averaged for each animal.

Preparation of Drugs

The drugs used in this study were milnacipran, citalopram, desipramine (provided by Asahi Kasei Corporation, Osaka, Japan), NMDA, AMPA, CNQX, bicuculline, strychnine, WAY100635 maleate salt (Sigma-Aldrich, St. Louis, MO) and tetrodotoxin (TTX), yohimbine hydrochloride (Wako, Osaka, Japan). Citalopram and bicuculline were first dissolved in dimethyl sulphoxide (DMSO) at 1000 times the concentrations to be used. The other drugs were first dissolved in distilled water at 1000 times the concentrations to be used, and then these drugs were diluted to the final concentration in Krebs solution immediately before use.

Statistical Analysis

Data are expressed as means \pm SEM. Statistical significance was determined as $P < 0.05$ using either the Student's paired t -test or a two-way analysis of variance (ANOVA), followed by a Student-Newman-Keuls post hoc test for multiple comparisons.


Results

Milnacipran Suppresses NMDA-Induced Thermal Hyperalgesia


Intrathecal injection of saline or each antidepressant alone did not alter PWL (data not shown). Intrathecal injection of 1 μ g NMDA 30 min after saline injection produced thermal hyperalgesia 30 min after the NMDA injection compared with the pre-value (9.87 ± 0.89 to 6.13 ± 0.89 s, $P < 0.01$). Mechanical allodynia was not induced by intrathecal injection of NMDA. Although the hyperalgesia induced by intrathecal administration of NMDA is a rapid and transient in a tail-flick test [19], the reduction in PWL continued up to 120 min after NMDA injection in the present study (saline group, Figure 1). Intrathecal injection of milnacipran followed by intrathecal injection of NMDA suppressed thermal hyperalgesia in a dose-dependent manner at doses from 0.01 to 0.1 μ mol ($P < 0.05$ by two-way ANOVA, Figure 1A). In contrast, citalopram (0.1 μ mol) did not inhibit thermal hyperalgesia during the testing period ($P = 0.25$ by two-way ANOVA; Figure 1B). Similarly, desipramine (0.1 μ mol) did not suppress NMDA-induced thermal hyperalgesia ($P = 0.62$ by two-way ANOVA, Figure 1C).

Milnacipran Inhibits NMDA, but not AMPA Receptor-Mediated Responses in Dorsal Horn Neurons

To study the effects of milnacipran on excitatory synaptic transmission, whole-cell patch-clamp recordings were made from rat lamina II neurons. Milnacipran did not alter the level of holding current required to maintain neurons at -40 mV and -70 mV, respectively. Milnacipran had no effect on the amplitude ($102 \pm 5\%$ of control, $n = 4$, $P = 0.81$) and frequency ($97 \pm 6\%$ of



control, $n = 4$, $P = 0.62$) of spontaneous EPSCs at -70 mV. Exogenous application of NMDA ($100 \mu\text{M}$, 30 s, at -40 mV) elicited an inward current in neurons (Figure 2A, C), reflecting the activation of NMDA receptors. To confirm that NMDA-induced currents were postsynaptic phenomena, we examined the currents in the presence of TTX to remove any possible



influence of the NMDA receptors on presynaptic neurons. Since TTX (0.5 μ M) did not affect the amplitudes of NMDA-induced currents (103 ± 4 % of control, $n=5$; $P=0.84$; data not shown), the currents were exclusively postsynaptic. Therefore, the following experiments were done in the absence of TTX. Pre-application of milnacipran (100 μ M) for 3 min reduced the amplitudes of NMDA-induced currents to 56 ± 3 % ($n=10$, $P<0.01$; Fig. 2A, B) of the control values. These effects of milnacipran were reversible, and the amplitudes of currents recovered to the control values within 5–10 min (Figure 2A). At a lower concentration of 10 or 30 μ M, milnacipran also decreased NMDA-induced currents (80 ± 4 % of the control, $n=5$, $P<0.01$; 71 ± 8 % of the control, $n=6$, $P<0.05$, respectively; Figure 2B). To elucidate whether

the observed effects were specific for milnacipran, we examined other antidepressants; i.e., citalopram and desipramine. Pre-application of citalopram (100 μ M) for 3 min failed to reduce the amplitudes of NMDA-induced currents (100 ± 5 % of control, $n=5$, $P=0.96$; Figure 2C, D). Pre-application of desipramine (30 μ M) for 3 min also failed to reduce the amplitudes of NMDA-induced currents (98 ± 4 % of control, $n=5$, $P=0.57$; Figure 2C, D). In contrast, milnacipran (100 μ M) had no effect on the amplitudes of exogenously applied AMPA (10 μ M, 30 s, at -70 mV)-induced currents (98 ± 5 % of control, $n=7$, $P=0.63$; Figure 3A). These results suggest that milnacipran, but not the other antidepressants, inhibited NMDA, but not AMPA receptor-mediated responses in lamina II neurons.




Figure 3 (A) Milnacipran does not inhibit AMPA receptor-mediated responses in dorsal horn neurons. Milnacipran did not inhibit AMPA-induced currents. **(B)** Representative traces of dorsal root stimulation evoked monosynaptic NMDA- and AMPA-mediated EPSCs. NMDA-mediated EPSCs were recorded at $+40$ mV. AMPA-mediated EPSCs were recorded at -70 mV. Milnacipran reversibly decreased the amplitudes of NMDA-, but not AMPA-mediated EPSCs. **(C)** Comparison of the NMDA- and AMPA-mediated EPSCs in control and in the presence of milnacipran. * $P<0.01$.

We next tested the effects of milnacipran on the amplitudes of dorsal root stimulation evoked EPSCs. Milnacipran (100 μ M) inhibited the amplitudes of monosynaptic NMDA-mediated EPSCs to 64 ± 5 % ($n = 4$, $P < 0.01$; Figure 3B, C) of the control values. In contrast, milnacipran (100 μ M) did not inhibit the amplitudes of monosynaptic AMPA receptor-mediated EPSCs (97 ± 5 % of control, $n = 5$, $P = 0.62$; Figure 3B, C). To exclude the possibility that milnacipran inhibited the amplitudes of NMDA-mediated EPSCs as a result of the blockade of 5-HT and NA reuptake, we examined the NMDA-mediated EPSCs in the presence of 5-HT_{1A} receptor antagonist, WAY100635 (10 μ M), and α 2 receptor antagonist, yohimbine (1 μ M). The concentrations of WAY100635 and yohimbine at the concentrations used here are sufficiently high to block the 5-HT_{1A} [20] and α 2 receptors [21], respectively. However, milnacipran also inhibited the amplitudes of NMDA-mediated EPSCs to 69 ± 8 % ($n = 3$, $P < 0.01$) of the control values.

Effects of Antidepressants on ERK Activation in the Spinal Cord

The slice preparation offers a reliable condition under which to study pERK expression [22,23]. In addition, multiple slices can be prepared from each spinal cord segment, and milnacipran can be applied in a known condition *in vitro*. The slices were perfused for more than 3 h before stimulation to reduce possible pERK background caused by slice preparation [22,23]. There were very few pERK-positive neurons noted in control spinal cord slices (Figure 4).

In the present research, pERK-positive neurons were occasionally observed in the control dorsal horn. Bath application of NMDA (100 μ M) for 5 min produced

activation of pERK in the superficial dorsal horn neurons. Distribution of NMDA-evoked pERK expression was similar to that of previous observations [22]. The number of pERK-positive neurons significantly increased in the NMDA-treated group ($P < 0.01$ vs. control, Figure 4). The number of pERK-positive neurons in the group treated with milnacipran (100 μ M), but not with citalopram (100 μ M) or desipramine (30 μ M) followed by NMDA was significantly lower compared with the NMDA-alone group ($P < 0.05$, Figure 4).

Discussion

We demonstrated that intrathecal administration of milnacipran, but not citalopram or desipramine mediated an inhibition of NMDA-induced thermal hyperalgesia. Moreover, we documented the inhibition of NMDA-mediated currents by milnacipran, but not by citalopram or desipramine in spinal lamina II. We also demonstrated that activation of pERK induced by NMDA was significantly suppressed by milnacipran in dorsal horn neurons. Taken together, these findings indicate that milnacipran has a direct antinociceptive effect in the spinal cord through its modulation of NMDA receptors.

Some investigators have reported interactions between TCAs and NMDA receptors in nociceptive transmission. Eisenach and Gebhart [5] reported that intrathecal administration of amitriptyline reversed thermal hyperalgesia via NMDA receptor antagonism in a rat model of inflammation. Kawamata et al. [24] also using the rat model of inflammation, reported that intrathecally injected desipramine produced analgesia unrelated to NA reuptake inhibition. These reports suggest that TCAs exert a direct inhibitory effect on NMDA receptors to produce analgesia in the spinal cord. TCAs may




Figure 4 Effects of antidepressants on ERK activation in the spinal cord. ERK activation was significantly suppressed by simultaneous treatment with milnacipran compared with the NMDA-alone group, but not with citalopram or desipramine. Arrow indicates the pERK-positive neurons. NS = not significant.

not only inhibit 5-HT and NA reuptake, but could also interact with various receptors. It has been shown that TCAs block sodium channels [25,26] as well as voltage-dependent calcium channels [27,28] and that TCAs inhibit adenosine reuptake [29]. Further, most TCAs have affinity for opioid [30], NA, 5-HT, histamine, and muscarinic acetylcholine receptors [31]. Therefore, these various mechanisms of action of TCAs might contribute to antinociceptive effects in some kinds of chronic pain models.

Desipramine did not suppress NMDA-induced thermal hyperalgesia in the present study. However, Hwang and Wilcox [6] reported that desipramine was antinociceptive in three nociceptive tests, tail-flick test, intrathecal substance P-induced behavioral test and intradermal hyperalgesic saline-induced behavioral test. Moreover, they indicated that the analgesic effect by desipramine probably involves blockade of monoamine reuptake. This discrepancy between our result and that of previous study is likely to be due to different underlying mechanisms of desipramine.

Although there is some evidence that TCAs block NMDA receptor-mediated responses, the site of action is controversial. Based on radioligand binding studies, Reynolds and Miller [32] have suggested that TCAs act at the Zn^{2+} recognition site on the NMDA receptor. Sills and Loo [33] have reported that TCAs bind with higher affinity to the phencyclidine binding site on the NMDA receptor. Moreover, Sernagor et al. [34] have reported that desipramine blocked NMDA-induced currents in hippocampal neurons by acting on the open channel. In contrast to TCAs, milnacipran has no relevant affinity for any other receptors, including α -adrenergic, 5-HT, histamine, muscarinic acetylcholine, opioid, or NMDA receptors [35]. However, Shuto et al. [36] reported that milnacipran ($IC_{50} = 6.3 \mu M$), is a class of noncompetitive NMDA receptor antagonist, although the binding affinity of milnacipran for the NMDA receptor is not strong. Although there is no evidence that milnacipran binds to the NMDA receptor, it is assumed that it may act at the recognition site for Zn^{2+} or at the phencyclidine binding site of the NMDA receptor. Further study is required to clarify this point.

Milnacipran inhibits the presynaptic reuptake of monoamines, 5-HT and NA with an IC_{50} of 100 to 200 nM, respectively in the brain [37]. However, in this study, milnacipran has the antagonistic effect on NMDA-mediated responses in the spinal cord at a concentration of 10–100 μM . Some previous studies [32,36,38] indicate that TCAs also inhibit the NMDA receptors in the brain at a concentration of 10–100 μM . Therefore, it is likely that the concentration of inhibiting the NMDA receptors by milnacipran is higher than that of inhibiting reuptake of the monoamines.

In the present study, milnacipran reduced the amplitudes of exogenously applied NMDA-induced currents in lamina II neurons. Moreover, milnacipran inhibited the amplitudes of dorsal root stimulation evoked NMDA-mediated EPSCs. There are no differences in the degree of depression by milnacipran between NMDA induced-current and dorsal root stimulation evoked NMDA-mediated EPSCs. These results suggest that synaptic and extra synaptic NMDA receptors in dorsal horn neurons are similarly modulated by milnacipran. Moreover, milnacipran inhibited the amplitudes of NMDA-mediated EPSCs in the presence of 5-HT antagonist and $\alpha 2$ receptor antagonist. Therefore, it is unlikely that an antagonistic effect of milnacipran on NMDA receptors is mediated by 5-HT or NA receptors.

We observed that the highest dose of intrathecal milnacipran completely reversed thermal hyperalgesia induced by NMDA. In contrast, desipramine and citalopram did not produce any inhibitory effect, although the concentrations of desipramine and citalopram in this study are sufficiently high to inhibit reuptake of 5-HT or NA, respectively. These results suggest that increases in 5-HT or NA alone in the spinal cord have no effect on NMDA-induced thermal hyperalgesia. The antagonistic action of milnacipran for NMDA receptors may produce additional effects for some types of chronic pain. Previous studies demonstrated that intrathecal administration of milnacipran produced antiallodynic effects in rats with peripheral nerve injury [13,39,40], and that the effect was not completely reversed by an $\alpha 2$ receptor antagonist or a 5-HT receptor antagonist [13]. Therefore, it is conceivable that an antagonistic action for NMDA receptors contributes to the antiallodynic effect of milnacipran. Further studies are required to clarify the molecular mechanisms underlying the inhibitory effect of milnacipran on NMDA-mediated responses in dorsal horn neurons. In addition, to elucidate whether the observed effects were specific for milnacipran, further investigations using another SNRI are necessary to resolve this question.

ERK activation is detected in the spinal dorsal horn neurons after stimulation of nociceptive primary afferents and contributes to the development of central sensitization [22]. Activation of the NMDA receptor is partly involved in ERK induction following nociceptive stimulation [22]. Analgesic drugs such as local anesthetics [41], opioids, or cannabinoids [42] inhibit ERK induction in the spinal cord. In the present study, we demonstrated that milnacipran inhibited ERK induction following application of NMDA in the spinal cord. This result is consistent with our behavioral data showing that milnacipran attenuated thermal hyperalgesia following intrathecal NMDA injection. Our electrophysiological data clearly indicate that the direct inhibitory effect of milnacipran on NMDA-mediated current underlies these phenomena.

We show in the present study that milnacipran has an antagonistic effect of NMDA receptors in the spinal cord when administered intrathecally. However, it is not clear whether milnacipran has the similar effect in other central and peripheral tissue such as brain and skin. Moreover, NMDA receptor antagonists such as ketamine and phencyclidine have psychotomimetic or anti-depressant properties in humans when administered systemically. There have been no studies demonstrating that milnacipran has these similar properties in humans. Therefore, further study is required to clarify this point.

NMDA glutamate receptors are one of the major receptor channel types mediating rapid excitatory neurotransmission in the central nervous system. This receptor is composed of subunits from at least two families, NR1 and NR2. The NR1 subunit is essential for the function of NMDA receptors and is ubiquitously expressed in most neurons. The functional properties of NMDA receptors are determined by the NR2 subunit composition (NR2A–2D). Previous reports have demonstrated that desipramine inhibited NMDA-evoked responses in hippocampal neurons [32,43], but not in the present study. The respective NR2 subunits show different expression patterns in various regions of the brain and spinal cord. Whereas NR2A and NR2B subunits are prominent in the hippocampus [44], these subunits are not identified in spinal dorsal horn neurons [45]. This different composition of NMDA receptors may underlie the variability among tissues in the effects of desipramine on NMDA receptor-responses.

NMDA glutamate receptors also play a key role in central sensitization in chronic pain [3,4]. The pursuit of an NMDA receptor antagonist for the relief of chronic pain dates from the late 1980s when it was shown that NMDA antagonists inhibit the “wind-up” response [46,47]. The central sensitization that occurs in the spinal dorsal horn is held to be an important event in the pathway leading to neuropathic pain. For this reason, ketamine is currently widely applied in the treatment of neuropathic pain and for some kinds of chronic pain, including fibromyalgia [48]. Although there is little evidence that NMDA receptor antagonism is involved in the antinociceptive effect of milnacipran, milnacipran has been widely used in patients with fibromyalgia and has provided them with pain relief [49,50]. These effects of milnacipran for chronic pain have been previously discussed from the perspective of one aspect: balanced inhibition of the reuptake of 5-HT and NA. However, in the present study, our data suggest that milnacipran not only inhibits 5-HT and NA reuptake, but also acts as an NMDA receptor antagonist. Milnacipran may have the potential to prevent or suppress chronic pain related to central sensitization, especially when injected into intrathecal or epidural space.

Conclusions

In conclusion, the present study suggests that milnacipran, but not citalopram and desipramine, inhibits NMDA-induced glutamatergic transmission in rat dorsal horn neurons. These results suggest that not only is inhibition of 5-HT and NA reuptake an important factor for the analgesia induced by milnacipran in the spinal cord, but that inhibition of glutamate NMDA receptors is critical as well.

Competing interests

The authors declare that they have no competing interests.

Acknowledgements

This work was supported by a Grant-In-Aid for Scientific Research (grant number 20390414, 09 F09359, TK; 20591823, HO; 20591841, FA) from the Ministry of Education, Culture, Sports, Science and Technology and by a Grant-In-Aid for Scientific Research from Ministry of Health, Labour and Welfare, Tokyo, Japan.

Author details

¹Department of Anesthesiology, Niigata University Graduate School of Medical and Dental Sciences, 757 Asahimachi, Chuo-ku Niigata 951-8510, Japan. ²Department of Anesthesiology, Gunma University Graduate School of Medicine. ³Department of Anesthesiology, Kyoto Prefectural University of Medicine. ⁴Pain Mechanism Research Group.

Authors' contributions

MK and HO performed the animal surgery and behavioral test. TK carried out *in vitro* patch-clamp recordings. MS and FA conducted immunohistochemistry. SS performed the statistical analysis. TK, HO and FA drafted the manuscript. All authors read and approved the final manuscript.

Received: 2 February 2012 Accepted: 19 June 2012

Published: 19 June 2012

References

1. Dworkin RH, O'Connor AB, Backonja M, Farrar JT, Finnerup NB, Jensen TS, Kalso EA, Loeser JD, Miaskowski C, Nurmikko TJ, et al: **Pharmacologic management of neuropathic pain: evidence-based recommendations.** *Pain* 2007, **132**:237–251.
2. Sindrup SH, Otto M, Finnerup NB, Jensen TS: **Antidepressants in the treatment of neuropathic pain.** *Basic Clin Pharmacol Toxicol* 2005, **96**:399–409.
3. Ji RR, Kohno T, Moore KA, Woolf CJ: **Central sensitization and LTP: do pain and memory share similar mechanisms?** *Trends Neurosci* 2003, **26**:696–705.
4. South SM, Kohno T, Kaspar BK, Hegarty D, Vissel B, Drake CT, Ohata M, Jenab S, Sailer AW, Malkmus S, et al: **A conditional deletion of the NR1 subunit of the NMDA receptor in adult spinal cord dorsal horn reduces NMDA currents and injury-induced pain.** *J Neurosci* 2003, **23**:5031–5040.
5. Eisenach JC, Gebhart GF: **Intrathecal amitriptyline acts as an N-methyl-D-aspartate receptor antagonist in the presence of inflammatory hyperalgesia in rats.** *Anesthesiology* 1995, **83**:1046–1054.
6. Hwang AS, Wilcox GL: **Analgesic properties of intrathecally administered heterocyclic antidepressants.** *Pain* 1987, **28**:343–355.
7. Kumazawa T, Perl ER: **Excitation of marginal and substantia gelatinosa neurons in the primate spinal cord: indications of their place in dorsal horn functional organization.** *J Comp Neurol* 1978, **177**:417–434.
8. Yoshimura M, Nishi S: **Blind patch-clamp recordings from substantia gelatinosa neurons in adult rat spinal cord slices: pharmacological properties of synaptic currents.** *Neuroscience* 1993, **53**:519–526.
9. Yaksh TL, Rudy TA: **Chronic catheterization of the spinal subarachnoid space.** *Physiol Behav* 1976, **17**:1031–1036.
10. Hargreaves K, Dubner R, Brown F, Flores C, Joris J: **A new and sensitive method for measuring thermal nociception in cutaneous hyperalgesia.** *Pain* 1988, **32**:77–88.
11. Malmberg AB, Yaksh TL: **Hyperalgesia mediated by spinal glutamate or substance P receptor blocked by spinal cyclooxygenase inhibition.** *Science* 1992, **257**:1276–1279.

12. Obata H, Kimura M, Nakajima K, Tobe M, Nishikawa K, Saito S: Monoamine-dependent, opioid-independent antihypersensitivity effects of intrathecally administered milnacipran, a serotonin noradrenaline reuptake inhibitor, in a postoperative pain model in rats. *J Pharmacol Exp Ther* 2010, **334**:1059–1065.
13. Obata H, Saito S, Koizuka S, Nishikawa K, Goto F: The monoamine-mediated antialloodynic effects of intrathecally administered milnacipran, a serotonin noradrenaline reuptake inhibitor, in a rat model of neuropathic pain. *Anesth Analg* 2005, **100**:1406–1410. table of contents.
14. Otsuka N, Kiuchi Y, Yokogawa F, Masuda Y, Oguchi K, Hosoyamada A: Antinociceptive efficacy of antidepressants: assessment of five antidepressants and four monoamine receptors in rats. *J Anesth* 2001, **15**:154–158.
15. Yokogawa F, Kiuchi Y, Ishikawa Y, Otsuka N, Masuda Y, Oguchi K, Hosoyamada A: An investigation of monoamine receptors involved in antinociceptive effects of antidepressants. *Anesth Analg* 2002, **95**:163–168. table of contents.
16. Kohno T, Wakai A, Ataka T, Ikoma M, Yamakura T, Baba H: Actions of midazolam on excitatory transmission in dorsal horn neurons of adult rat spinal cord. *Anesthesiology* 2006, **104**:338–343.
17. Kohno T, Wang H, Amaya F, Brenner GJ, Cheng JK, Ji RR, Woolf CJ: Bradykinin enhances AMPA and NMDA receptor activity in spinal cord dorsal horn neurons by activating multiple kinases to produce pain hypersensitivity. *J Neurosci* 2008, **28**:4533–4540.
18. Amaya F, Samad TA, Barrett L, Broom DC, Woolf CJ: Perianglionic inflammation elicits a distally radiating pain hypersensitivity by promoting COX-2 induction in the dorsal root ganglion. *Pain* 2009, **142**:59–67.
19. Meller ST, Dykstra C, Gebhart GF: Acute thermal hyperalgesia in the rat is produced by activation of N-methyl-D-aspartate receptors and protein kinase C and production of nitric oxide. *Neuroscience* 1996, **71**:327–335.
20. Ito A, Kumamoto E, Takeda M, Shibata K, Sagai H, Yoshimura M: Mechanisms for ovariectomy-induced hyperalgesia and its relief by calcitonin: participation of 5-HT1A-like receptor on C-afferent terminals in substantia gelatinosa of the rat spinal cord. *J Neurosci* 2000, **20**:6302–6308.
21. Kawasaki Y, Kumamoto E, Furue H, Yoshimura M: Alpha 2 adrenoceptor-mediated presynaptic inhibition of primary afferent glutamatergic transmission in rat substantia gelatinosa neurons. *Anesthesiology* 2003, **98**:682–689.
22. Kawasaki Y, Kohno T, Zhuang ZY, Brenner GJ, Wang H, Van Der Meer C, Befort K, Woolf CJ, Ji RR: Ionotropic and metabotropic receptors, protein kinase A, protein kinase C, and Src contribute to C-fiber-induced ERK activation and cAMP response element-binding protein phosphorylation in dorsal horn neurons, leading to central sensitization. *J Neurosci* 2004, **24**:8310–8321.
23. Ji RR, Baba H, Brenner GJ, Woolf CJ: Nociceptive-specific activation of ERK in spinal neurons contributes to pain hypersensitivity. *Nat Neurosci* 1999, **2**:1114–1119.
24. Kawamata T, Omote K, Kawamata M, Namiki A: Analgesic effect of intrathecal desipramine on carrageenan-induced thermal hyperalgesia in the rat. *Br J Anaesth* 1999, **83**:449–452.
25. Pancrazio JJ, Kamatchi GL, Roscoe AK, Lynch C 3rd: Inhibition of neuronal Na⁺ channels by antidepressant drugs. *J Pharmacol Exp Ther* 1998, **284**:208–214.
26. Dick IE, Brochu RM, Purohit Y, Kaczorowski GJ, Martin WJ, Priest BT: Sodium channel blockade may contribute to the analgesic efficacy of antidepressants. *J Pain* 2007, **8**:315–324.
27. Lavoie PA, Beauchamp G, Elie R: Tricyclic antidepressants inhibit voltage-dependent calcium channels and Na⁽⁺⁾-Ca²⁺ exchange in rat brain cortex synaptosomes. *Can J Physiol Pharmacol* 1990, **68**:1414–1418.
28. Shimizu M, Nishida A, Yamawaki S: Antidepressants inhibit spontaneous oscillations of intracellular Ca²⁺ concentration in rat cortical cultured neurons. *Neurosci Lett* 1992, **146**:101–104.
29. Phillis JW, Wu PH: The effect of various centrally active drugs on adenosine uptake by the central nervous system. *Comp Biochem Physiol C* 1982, **72**:179–187.
30. Isenberg KE, Cicero TJ: Possible involvement of opiate receptors in the pharmacological profiles of antidepressant compounds. *Eur J Pharmacol* 1984, **103**:57–63.
31. Hall H, Ogren SO: Effects of antidepressant drugs on different receptors in the brain. *Eur J Pharmacol* 1981, **70**:393–407.
32. Reynolds IJ, Miller RJ: Tricyclic antidepressants block N-methyl-D-aspartate receptors: similarities to the action of zinc. *Br J Pharmacol* 1988, **95**:95–102.
33. Sills MA, Loo PS: Tricyclic antidepressants and dextromethorphan bind with higher affinity to the phencyclidine receptor in the absence of magnesium and L-glutamate. *Mol Pharmacol* 1989, **36**:160–165.
34. Sernagor E, Kuhn D, Vyklícky L Jr, Mayer ML: Open channel block of NMDA receptor responses evoked by tricyclic antidepressants. *Neuron* 1989, **2**:1221–1227.
35. Mochizuki D, Tsujita R, Yamada S, Kawasaki K, Otsuka Y, Hashimoto S, Hattori T, Kitamura Y, Miki N: Neurochemical and behavioural characterization of milnacipran, a serotonin and noradrenaline reuptake inhibitor in rats. *Psychopharmacology (Berl)* 2002, **162**:323–332.
36. Shuto S, Takada H, Mochizuki D, Tsujita R, Hase Y, Ono S, Shibuya N, Matsuda A: (+/-)-(Z)-2-(aminomethyl)-1-phenylcyclopropanecarboxamide derivatives as a new prototype of NMDA receptor antagonists. *J Med Chem* 1995, **38**:2964–2968.
37. Moret C, Charveron M, Finberg JP, Couzinier JP, Briley M: Biochemical profile of midalcipran (F 2207), 1-phenyl-1-diethyl-aminocarbonyl-2-aminomethyl-cyclopropane (Z) hydrochloride, a potential fourth generation antidepressant drug. *Neuropharmacology* 1985, **24**:1211–1219.
38. Watanabe Y, Saito H, Abe K: Tricyclic antidepressants block NMDA receptor-mediated synaptic responses and induction of long-term potentiation in rat hippocampal slices. *Neuropharmacology* 1993, **32**:479–486.
39. Shin SW, Eisenach JC: Peripheral nerve injury sensitizes the response to visceral distension but not its inhibition by the antidepressant milnacipran. *Anesthesiology* 2004, **100**:671–675.
40. King T, Rao S, Vanderah T, Chen Q, Vardanyan A, Porreca F: Differential blockade of nerve injury-induced shift in weight bearing and thermal and tactile hypersensitivity by milnacipran. *J Pain* 2006, **7**:513–520.
41. Yanagidate F, Strichartz GR: Bupivacaine inhibits activation of neuronal spinal extracellular receptor-activated kinase through selective effects on ionotropic receptors. *Anesthesiology* 2006, **104**:805–814.
42. Kawasaki Y, Kohno T, Ji RR: Different effects of opioid and cannabinoid receptor agonists on C-fiber-induced extracellular signal-regulated kinase activation in dorsal horn neurons in normal and spinal nerve-ligated rats. *J Pharmacol Exp Ther* 2006, **316**:601–607.
43. White G, Lovinger DM, Peoples RW, Weight FF: Inhibition of N-methyl-D-aspartate activated ion current by desmethylimipramine. *Brain Res* 1990, **537**:337–339.
44. Monyer H, Burnashev N, Laurie DJ, Sakmann B, Seeburg PH: Developmental and regional expression in the rat brain and functional properties of four NMDA receptors. *Neuron* 1994, **12**:529–540.
45. Tolle TR, Berthele A, Zieglgansberger W, Seeburg PH, Wisden W: The differential expression of 16 NMDA and non-NMDA receptor subunits in the rat spinal cord and in periaqueductal gray. *J Neurosci* 1993, **13**:5009–5028.
46. Davies SN, Lodge D: Evidence for involvement of N-methylaspartate receptors in 'wind-up' of class 2 neurones in the dorsal horn of the rat. *Brain Res* 1987, **424**:402–406.
47. Dickenson AH, Sullivan AF: Evidence for a role of the NMDA receptor in the frequency dependent potentiation of deep rat dorsal horn nociceptive neurones following C fibre stimulation. *Neuropharmacology* 1987, **26**:1235–1238.
48. Domino EF: Taming the ketamine tiger. 1965. *Anesthesiology* 2010, **113**:678–684.
49. Gendreau RM, Thorn MD, Gendreau JF, Kranzler JD, Ribeiro S, Gracely RH, Williams DA, Mease PJ, McLean SA, Clauw DJ: Efficacy of milnacipran in patients with fibromyalgia. *J Rheumatol* 2005, **32**:1975–1985.
50. Mease PJ, Clauw DJ, Gendreau RM, Rao SG, Kranzler J, Chen W, Palmer RH: The efficacy and safety of milnacipran for treatment of fibromyalgia: a randomized, double-blind, placebo-controlled trial. *J Rheumatol* 2009, **36**:398–409.

doi:10.1186/1744-8069-8-45

Cite this article as: Kohno et al.: Milnacipran inhibits glutamatergic N-Methyl-D-Aspartate receptor activity in Spinal Dorsal Horn Neurons. *Molecular Pain* 2012 **8**:45.