

REVIEW

Open Access

Roles of phosphatase 2A in nociceptive signal processing

Yun Wang^{1*}, Yongzhong Lei², Li Fang², Yonggao Mu³, Jing Wu⁴ and Xuan Zhang^{2*}

Abstract

Multiple protein kinases affect the responses of dorsal horn neurons through phosphorylation of synaptic receptors and proteins involved in intracellular signal transduction pathways, and the consequences of this modulation may be spinal central sensitization. In contrast, the phosphatases catalyze an opposing reaction of de-phosphorylation, which may also modulate the functions of crucial proteins in signaling nociception. This is an important mechanism in the regulation of intracellular signal transduction pathways in nociceptive neurons. Accumulated evidence has shown that phosphatase 2A (PP2A), a serine/threonine specific phosphatase, is implicated in synaptic plasticity of the central nervous system and central sensitization of nociception. Therefore, targeting protein phosphatase 2A may provide an effective and novel strategy for the treatment of clinical pain. This review will characterize the structure and functional regulation of neuronal PP2A and bring together recent advances on the modulation of PP2A in targeted downstream substrates and relevant multiple nociceptive signaling molecules.

Introduction

Intracellular signal transduction pathways play a pivotal role in the maintenance of biological processes such as cell growth, proliferation, survival, and metabolism in all cells and tissues. It has been demonstrated that a variety of intracellular signal transduction pathways are involved in the physiological or patho-physiological responses to noxious stimuli [1-3]. The opposing reactions of phosphorylation and de-phosphorylation of critical cellular proteins are decisive to such pathways [4,5]. Protein kinases and phosphatases catalyze protein phosphorylation and de-phosphorylation reactions, respectively. While in the past, much attention has been paid to the regulation of protein kinases, it is now apparent that protein phosphatases are highly regulated enzymes that play an equally important role in the control of protein phosphorylation. Accumulated evidence has shown that protein kinases are widely implicated in pain modulation. Several protein kinases affect the responses of spinal cord dorsal horn neurons through phosphorylation of synaptic receptors and proteins involved in intracellular signal transduction

pathways, and the consequences of this modulation can regulate the process of central sensitization [2,6-10]. However, much less is known about the role of their counterparts, protein phosphatases, in nociception. Recent studies have provided evidence that a member of protein phosphatase family, protein phosphatase 2A (PP2A), is involved in synaptic plasticity in the central nervous system (CNS) or central sensitization of pain, suggesting a new promising molecular target for pain control [11-15].

Serine/threonine specific phosphatase is one of major classes of protein phosphatases that catalyze the de-phosphorylation of serine and threonine residues. According to their biological characteristics, sensitivities to specific inhibitors and substrates, serine/threonine specific phosphatase can be divided into four major subtypes, PP1, PP2A, PP2B and PP2C [16]. Among this family members, PP2C belongs to a separate gene family since it has a distinct structure from the others, whereas PP1, PP2A and PP2B have similar primary amino acid sequences. There are other serine/threonine phosphatases identified as well, including PP4, PP5, PP6 and PP7. Unlike PP1 and PP2A, the *in vitro* basic activities of PP4, PP5, PP6, and PP7 are extremely low. Of these subtypes, PP2A is the most abundant serine/threonine protein phosphatase in mammalian cells and is expressed at higher levels in the CNS [17]. This review will characterize the structure and functional regulations of PP2A and

* Correspondence: sincerewy@yahoo.com; Xuanczhang@gmail.com

¹Department of Anesthesiology, Beijing Chaoyang Hospital, Capital Medical University, Beijing 100020, China

²Department of Neuroscience and Cell Biology, The University of Texas Medical Branch, Galveston, TX 77555-0517, USA

Full list of author information is available at the end of the article

highlight recent advances in the involvement of PP2A in de-phosphorylation of specific downstream substrates and nociceptive signal processing.

The structure and localization of PP2A

PP2A is a major serine/threonine protein phosphatase in mammalian cells and has been implicated in the control of numerous biological processes including development, cell growth, differentiation, and apoptosis. It accounts for up to 1% of all cellular proteins and, together with PP1, accounts for 90% of all serine/threonine phosphatase activity in most tissues and cells [18]. It predominantly exists in cells as a heterotrimeric holoenzyme, which consists of a 36 kDa catalytic subunit (PP2A-C), a 65 kDa structural subunit (PP2A-A) forming a core enzyme, and a variable regulatory subunit (PP2A-B), as illustrated in Figure 1 [19]. The A structure subunit recruits the C catalytic subunit to form the core dimer, which acts as a scaffold for B subunits of the enzyme. Four B subunit families have been identified (PR55 or B, PR61 or B', PR72/130 or B'' and PR93/PR110 or B'''). Different B subunits interact via the same or overlapping sites within the A subunit of the core dimer. The association of these B subunits with the core AC dimer is




Figure 1 Structure of protein phosphatase 2A holoenzymes. A is the structural subunit and scaffolding protein, and C is the catalytic subunit. B/B'/B''/B''' are the variable and regulatory subunits. PP2A predominantly exists in cells as a heterotrimeric holoenzyme, which consists of a 36 kDa catalytic subunit (PP2A-C), a 65kDa structural subunit (PP2A-A) forming a core enzyme, and a variable regulatory subunit (PP2A-B). The A structure subunit recruits the C catalytic subunit to form the core dimer, which acts as a scaffold for the C and B subunits of the enzyme. Four B subunit families including PR55 or B, PR61 or B', PR72/130 or B'' and PR93/PR110 or B''' interact via the same or overlapping sites within the A subunit of the core dimer. The PP2A holoenzyme's substrate specificity, enzymatic activity, and/or cellular localization can be modulated by the B regulatory subunit.

mutually exclusive. The PP2A holoenzyme's substrate specificity, enzymatic activity, and/or cellular localization can be modulated by the B regulatory subunit [20-23].

Each family of B subunits contains several isoforms that can bind the AC dimer in a mutually exclusive manner [24]. In mammalian cells, B or PR55 family (α , β , γ and δ) is expressed in a tissue-specific manner. PR55 α and PR55 δ are widely-distributed in different tissues, whereas PR55 β and PR55 γ are highly enriched in the brain. PR55 α is distributed primarily in the cell body and nucleus of Purkinje cells, whereas PR55 β is excluded from the nucleus and extends into dendrites. The B' family consists of five primary members of the PR61 (α , β , γ , δ and ϵ) that are mapped to the human chromosomes loci 1q41, 11q12, 3p21, 6p21.1, and 7p11.2, respectively. PR61 α , PR61 β and PR61 ϵ localize to the cytoplasm, whereas PR61 γ 1, PR61 γ 2, PR61 γ 3 are concentrated in the nucleus, and PR61 δ is found in both the nucleus and the cytoplasm. PR61 α and PR61 γ 1- γ 3 are enriched in heart and skeletal muscles. PR61 β and PR61 δ are expressed predominantly in the brain. For the B'' family, PR72 is expressed exclusively in the heart and skeletal muscles, whereas PR130 is expressed predominately in the heart and muscles. The PR93/PR110 comprises the B''' family. PR110 is localized to the post-synaptic densities of neuronal dendrites, whereas PR93 is nuclear located (Table 1) [16,22]. The spinal cord dorsal horn is an important area containing sensory neurons that has been implicated in major pain processing. Our previous immunofluorescent staining studies have shown that the neurons with PP2A expression are mainly distributed in the superficial layers of the dorsal horn, laminae I and II, and a few of the PP2A immunoreactive neurons are found in the ventral horn of the spinal cord as well (unpublished data).

The functional regulation of PP2A activity through post-translational modification

Previous studies have demonstrated that modulations of PP2A activity are due to the tyrosine or serine/threonine phosphorylation of not only the catalytic C subunit, but also the regulatory B subunit, especially those of the B' family [18,20]. The regulatory B subunits play key roles in controlling PP2A substrate specificity, cellular localization, and enzymatic activity [25]. Tyrosine kinases such as Src kinase, inhibit PP2A activity, and the PP2A assembly can be inhibited by the phosphorylation of the B56 subunit by extracellular signaling regulated kinase (ERK) [26,27]. Another post-translational modification of the catalytic C subunit, methylation, which occurs on the carboxy group of the C-terminal residue Leu³⁰⁹, is also involved in the alterations of PP2A's activity [28,29]. It has been shown that the methylation of PP2A-C may influence the affinity of the AC core dimer to the different B subunits [30,31]. For example, some B regulatory

Table 1 Tissue distribution and subcellular localization of PP2A subunits

Subunits	Molecule	KDa	Isoforms	Tissue distribution	Subcellular localization
Structure subunit	A	65	α and β	Ubiquitously expressed	
Catalytic subunit	C	36	α and β	Ubiquitously expressed. High levels in brain and heart.	
Regulatory subunit	B	55	B α	Wide-spread tissue distribution	Cytosolic fraction
			B β	Enriched in brain	Cytosolic fraction
			B δ	Wide-spread tissue distribution	
			B γ	Enriched in brain	Cytoskeletal fraction
			B' α	Widely expressed and abundant in heart and skeletal muscle	Cytoplasm
			B' β	Enriched in brain	Cytoplasm
			B' δ	Enriched in brain	Nucleus and cytoplasm
			B' γ	Widely expressed and abundant in heart and skeletal muscle	Nucleus
			B' ϵ		Cytoplasm
			B' PR72	Heart and skeletal muscle	
			B"PR130	Ubiquitously expressed and high levels in heart and muscle	
B"PR59	Testis, kidney, liver, brain, heart and lung				
B"PR48		nucleus			
B" PR93		nucleus			
B"PR110		Postsynaptic densities of neuronal dendrites			

subunits appear to bind to an AC dimer more efficiently when the catalytic C subunit has been methylated, whereas other B subunits prefer to bind an AC dimer with a demethylated C subunit [32]. The post-translational modification of PP2A has been implicated in the pathogenesis of Alzheimer's disease (AD), a neurodegenerative disorder with impaired synaptic plasticity [33-37]. The reduced methylation of PP2A C subunit at Leu³⁰⁹ and the increased phosphorylation of PP2A C subunit at Tyr³⁰⁷ may result in loss of enzymatic activity and tau hyperphosphorylation in Alzheimer's disease, indicating that PP2A is a putative target of therapeutic intervention [34]. How the post-translational modification of PP2A is regulated during nociception is still unclear and deserves further investigations.

Substrate molecules and signal transduction cascades regulated by PP2A in synaptic plasticity and central sensitization in response to nociceptive stimuli

It has been demonstrated that synaptic glutamate receptors play a critical role in synaptic plasticity, electrophysiologically characterized by long term potentiation (LTP) and long-term depression (LTD), in the central nervous system [38-41]. Previous studies from ours and other groups have demonstrated that central sensitization of pain may represent a spinal form of LTP, since there are close parallels in mechanisms important for LTP and central sensitization [1,38-40]. The functional activation of

post-synaptic glutamate receptors may influence a variety of intracellular signals, which may trigger cellular and molecular changes at transcriptional, translational, or post-translational levels. These changes contribute to the synaptic plasticity and central sensitization induced by peripheral noxious stimulation [3,6,42,43]. The phosphorylation and de-phosphorylation of synaptic glutamate receptors are important post-translational mechanisms in the modulation of synaptic strength. Strong noxious stimulation in the periphery may activate several protein kinases such as calcium/calmodulin dependent protein kinase II (CaMKII), cAMP-dependent protein kinase (PKA), protein kinase C (PKC), and protein kinase G (PKG), which play an important role in the phosphorylation of glutamate receptors in spinal nociceptive neurons [1,3,7,42-45]. The increased sensitivity of glutamate receptors through the phosphorylation regulated by protein kinases may contribute to the enhanced responsiveness of dorsal horn neurons during central sensitization [1,3,10]. In contrast, the protein phosphatase, PP2A may reverse these signals through the de-phosphorylation of glutamate receptors and several intracellular protein kinases, and therefore, blunt the central sensitization of pain [11-14].

PP2A is involved in the induction and maintenance of synaptic plasticity: electrophysiological evidence

The modification of protein phosphorylation is a critical element leading to the induction of synaptic plasticity.

For example, long-term potentiation is accompanied by increased glutamate receptor phosphorylation through various protein kinases and a concomitant decrease in protein phosphatase activity [46]. In contrast, a decrease in synaptic strength, LTD, has been shown to be dependent on glutamate receptor de-phosphorylation mediated by an increase in the activity of protein phosphatases, possibly PP1 and PP2A [47]. Thus, a coordination of kinase and phosphatase activities is crucial for the comprehensive modulation of synaptic plasticity.

The application of the PP1/PP2A inhibitor calyculin A, or the post-synaptic injection of microcystin, in hippocampal slices induces a rapid enhancement of synaptic transmission, particularly in the hippocampal tissue from aged rats [48]. It indicates that synaptic transmission can be actively and persistently regulated by protein phosphatases. It has been shown that PKA plays an important role in N-methyl-D-aspartic acid (NMDA) receptor-mediated plasticity in the hippocampus and spinal cord through the phosphorylation of GluN1 subunit of NMDA receptors. Both pharmacological and genetic inhibition of the cAMP/PKA cascade may inhibit the LTP in the hippocampal CA1 area [49,50]. The sensitivity of LTP (elicited by either multiple 100-Hz trains or prolonged 5-Hz stimulation) to PKA inhibition was eliminated by the prior incubation of hippocampal slices with a PP1/PP2A inhibitor, suggesting that the PKA pathway participates in LTP, which may include the activity-dependent suppression of PP1/PP2A activity [51,52]. It has been noted that the induction of LTP is associated with an inhibition of PP2A. The inhibition of PP2A activity was not only observed immediately after the induction of LTP, but was still detectable one hour after induction, indicating that persistently decreased PP2A activity may have a role in the maintenance of LTP [53,54]. Furthermore, the observed inhibition of PP2A showed a pattern of NMDA-receptors dependence. The auto-phosphorylation of CaMKII triggered by the activation of NMDA receptors is a cellular event critical to the induction of LTP. It is shown that the purified PP2A B α is a substrate for CaMKII phosphorylation, and this subunit is phosphorylated during the induction of LTP in the hippocampus [54]. Combined with these data, we may suggest that the inhibited activity of PP2A during LTP is associated with the phosphorylation of PP2A mediated by CaMKII triggered by the activation of NMDA receptor. As for the decrease in PP2A activity, this phosphorylation persisted for more than one hour after LTP induction. In view of this CaMKII-dependent decrease in PP2A activity, it is interesting that auto-phosphorylated CaMKII is much more readily de-phosphorylated by PP1 than by PP2A [54]. Thus, the CaMKII-dependent suppression of PP2A activity and prevention of the de-phosphorylation of CaMKII by PP2A might serve as key processing events necessary for the LTP maintenance

(Figure 2). Pi et al. [55] showed that the coupled PP2A and CaMKII switches lead to a tristable system in which the kinase activity is high in the LTP state, the PP2A activity is high in the LTD state, and neither activity is high in the basal state. These data provide an explanation for the inhibition of PP2A prevents LTD induction.

The auto-phosphorylation of CaMKII induced by tetanization is reported to be blocked by a PKA inhibitor, indicating a potential downstream substrate regulated by the PKA-dependent suppression of phosphatase activity. One possibility is that PP2A regulates LTP by competing with PKA for the regulation of specific phosphorylation sites, such as the GluN1 subunit of NMDA receptors. Although much more work needs to be done in this area, the data indicate an important involvement of the persistent down-regulation of PP2A activity in the maintenance of LTP. PP2A are also found to be involved in the development of LTD in the hippocampus. A number of studies showed that PP2A inhibitors disrupt NMDA receptor-dependent LTD at glutamatergic synapses in hippocampus [56,57].

PP2A is implicated in synaptic plasticity and pain central sensitization through regulating the function of NMDA receptors

NMDA receptors are nonselective cation channels critical for neuronal excitability and particularly for Ca²⁺-dependent modulation of synaptic plasticity in nociceptive processing. It was found that in the mouse nucleus accumbens, endogenous PP2A regulates NMDA receptor channel phosphorylation, activity and kinetics. In cultured hippocampal neurons, exogenous PP2A depressed the open probability of NMDA receptors [58]. Phosphatase may participate in the long-term changes in NMDA receptor function, such as LTP or LTD. Okadaic acid, which serves as a potent and specific inhibitor of the serine/threonine protein phosphatases 1 and 2A, may enhance the open probability of NMDA receptors [59,60]. It has been demonstrated that okadaic acid increases the NMDA and AMPA-kainate receptors-mediated currents in neurons of the hippocampus [61]. Intra-hippocampal micro-injection of okadaic acid induces the hyper-phosphorylation of the GluN2B subunit of the NMDA receptors [62]. Association of PP2A with NMDA receptors may result in an increase in the phosphatase activity of PP2A and the dephosphorylation of Serine⁸⁹⁷ of the NMDA receptor subunit NR1. On the other hand, the dissociation of PP2A from the complex and the reduction of PP2A activity can be caused by stimulation of NMDA receptors [63].

Using an animal model of inflammatory pain, our previous studies have demonstrated that PP2A plays a critical role in determining the excitability of nociceptive neurons in the spinal cord by modulating the phosphorylation state




Figure 2 Molecular substrates regulated by PP2A in synaptic plasticity and central sensitization of pain. The auto-phosphorylation of CaMKII triggered by the activation of NMDA receptors is an event critical to the induction of LTP. PP2A is a substrate for CaMKII phosphorylation and the phosphorylated PP2A may decrease the activity of PP2A during LTP. The CaMKII-dependent suppression of PP2A activity and prevention of the de-phosphorylation of CaMKII by PP2A may be necessary for the LTP maintenance. PP2A regulates LTP by competing with PKA for the regulation of specific phosphorylation sites, such as the GluN1 subunits of NMDA receptors. AMPA receptor GluA1 subunit has the two major phosphorylation sites: Ser⁸⁴⁵, which is phosphorylated by PKA, and Ser⁸³¹, which is phosphorylated by PKC. CaMKII was also found to phosphorylate both Ser⁸³¹ and Ser⁸⁴⁵ in GluA1, and contributes to the single-channel conductance of the receptor, and thus, possibly increases AMPA receptor conductance during LTP. The de-phosphorylation of Ser⁸⁴⁵ was blocked by the pretreatment with okadaic acid, indicating an involvement of PP1 and/or PP2A. The phosphorylation or de-phosphorylation of AMPA receptors is closely associated with the receptor trafficking. The GluA1 subunit is the important substrate of PP2A, indicating PP2A activity is a critical for AMPA receptor trafficking and might play an important role in AMPA receptor-mediated nociception. The transcription factor cAMP-response-element-binding protein (CREB) has been demonstrated to be involved in synaptic plasticity and gene transcription and PP2A is thought to be the main CREB phosphatase. Histone deacetylase (HDAC) may reverse the action of histone acetylase and block the gene transcription process through the chromatin remodeling. PP2A is responsible for the de-phosphorylation of class II HDACs and for the subsequent triggering nuclear localization and repression of target genes, while the phosphorylation-triggering cytoplasmic localization may lead to the activation of target genes.

of some critical proteins [13,14]. Infusion of a general inhibitor of PP2A, okadaic acid (OA), and a specific inhibitor, fostriecin into the subarachnoid space may significantly enhance the secondary mechanical hyperalgesia and allodynia following intradermal injection of capsaicin [14]. Further, we found that the up-regulated phosphorylation of both GluN1 and GluN2B subunits of NMDA receptors induced by capsaicin injection was significantly potentiated by the PP2A inhibitor without

affecting the GluN1 and GluN2B protein itself in the spinal cord dorsal horns. It suggests that PP2A may have a regulatory effect on central sensitization induced by noxious stimuli in the periphery by regulating the phosphorylation state of NMDA receptors [13]. Using an *in vivo* electrophysiological recording technique, our previous study has also shown that PP2A inhibitors significantly prolonged the responses of dorsal horn neurons to mechanical stimuli in anesthetized rats following intra-

dermal injection of capsaicin, indicating PP2A may be involved in determining the duration of capsaicin-induced central sensitization [11,12].

PP2A regulates the function of AMPA receptor through influencing the dephosphorylation and trafficking of AMPA receptor subunits

AMPA receptor has an important role in synaptic plasticity and central sensitization of pain [8,9,64-66]. AMPA receptors present unique functional regulations, such as subunits phosphorylation and membrane insertion or internalization. The intracellular C-terminal domains of AMPA receptor subunits allow subunit-specific regulation by phosphorylation. There are several protein phosphorylation sites located on the C-terminal region, which are working targets of PKA, PKC, and CaMKII [1,7]. Site-directed mutagenesis and phosphopeptide analysis identified two major phosphorylation sites on GluR1: Ser⁸⁴⁵, which is phosphorylated by PKA, and Ser⁸³¹, which is phosphorylated by PKC. The phosphorylation of Ser⁸⁴⁵ in GluA1 by PKA regulates the open-channel probability of AMPA receptors; whereas the phosphorylation of Ser⁸³¹ by PKC changes channel conductance. CaMKII was also found to phosphorylate both Ser⁸³¹ and Ser⁸⁴⁵ in GluA1 and contributes to the single-channel conductance of the receptor and possibly increases AMPA receptor conductance during LTP [65]. The phosphorylation of GluA2 plays an important role in the receptor clusters during synaptic plasticity and persistent painful stimulation [67]. It has been demonstrated that GluA2 may be phosphorylated on Ser⁸⁸⁰ by PKC *in vitro* and in transfected cells [68].

The development of phosphorylation-site-specific antibodies has greatly facilitated the study of the phosphorylation state of endogenous proteins regulated by kinases and phosphatases. Haganir's group has shown that the induction of LTD by prolonged, low-frequency stimulation led to a decrease in the phosphorylation of Ser⁸⁴⁵ but not Ser⁸³¹ of GluA1 subunits, as assessed by the Western blot analysis of hippocampal slices after stimulation [69]. The de-phosphorylation of Ser⁸⁴⁵ was blocked by pretreatment with okadaic acid, indicating an involvement of PP1 and/or PP2A (Figure 2). So, GluA1 is a critical substrate of protein phosphatases in LTD. Although a similar decrease in Ser⁸⁴⁵ phosphorylation was observed when LTD was induced chemically by the application of NMDA, this de-phosphorylation was not blocked by high concentrations of either okadaic acid or calyculin A [70]. This result indicates that different populations of GluA1 might be regulated by distinct phosphatases. Consistent with this idea, Strack and colleagues have observed a similar phenomenon in the case of another substrate, CaMKII. Whereas soluble CaMKII α is a PP2A substrate, translocation of CaMKII α to the postsynaptic density by auto-

phosphorylation converts it to a PP1 substrate [54]. Interestingly, it is noted that CaMKII activity in the postsynaptic density does not appear to correlate with synaptic insertion of GluA1 at C fiber synapses in inflammatory pain [41,71]. Thus, this strongly supports the notion that, even if PP1 is the major phosphatase in the postsynaptic density, cytosolic PP2A (or PP2B) may regulate the auto-phosphorylation of CaMKII that mediates synaptic AMPA receptor incorporation. However, the synaptic induction of LTD is associated with the de-phosphorylation of Ser⁸⁴⁵, but not Ser⁸³¹, and the induction of LTP in naive slices is observed to be associated with an increase in the phosphorylation of Ser⁸³¹, but not Ser⁸⁴⁵. LTP-inducing stimulation only elicited increased Ser⁸⁴⁵ phosphorylation when LTP was induced at previously depressed synapses. Conversely, at previously potentiated synapses, the administration of de-potentiating, low-frequency stimuli produced de-phosphorylation of Ser⁸³¹, but not Ser⁸⁴⁵ [69,72,73]. These studies show that the de-phosphorylation regulation by protein phosphatases in synaptic plasticity is associated with the functional status of the synapses. In spinal neurons, our group has shown that PKA mediates the phosphorylation of serine at the Ser⁸⁴⁵ site, and PKC targets the Ser⁸³¹ site following noxious stimulation [7]. Further, we have demonstrated that AMPA receptors showed enhanced responsiveness to nociceptive stimulation through this phosphorylation step during central sensitization [1]. However, much less is known about the de-phosphorylation regulation of GluA1 or GluA2 subunits by PP2A in different pain models.

The de-phosphorylation of ligand-gated channels regulates the channel properties. The accumulated evidence indicates that phosphatase activity also regulates the surface insertion or cluster of these neuronal receptors. Several studies have shown that NMDA-receptor-dependent LTD is associated with a post-synaptic silence of synapses, and that a post-synaptic interference with the endocytotic machinery hampers LTD. A series of studies now show that the rapid trafficking of AMPA receptors may occur in the hippocampal or spinal dorsal horn neurons after subunit phosphorylation [74,75]. The internalization of AMPA receptors can be blocked by inhibitors of protein phosphatase, indicating that protein phosphatase might have a regulatory role in LTD [76]. Currently, little is known about the role of PP2A in the trafficking of AMPA receptor subunits. It has been demonstrated that a membrane insertion of Ca²⁺-permeable AMPA receptors greatly contributes to the synaptic plasticity in hippocampus and central sensitization of pain in the spinal cord dorsal horns [8,77]. Haganir's group has also shown that the trafficking of AMPA receptors is regulated through the PKA phosphorylation of GluA1 subunits [78,79]. Another study indicated that the signaling pathway that drive the insertion of GluA1 subunits into the plasma membrane during

LTP *in vitro* require the activated CaMKII [80]. In spinal neurons, intrathecal application of a CaMKII inhibitor, KN-93, before the painful visceral stimulus, apparently inhibits the GluA1 accumulation in the plasma membrane fraction. Peripheral inflammation stimuli drive the phosphorylation and trafficking of AMPA receptor subunits in spinal cord dorsal horns [81,82]. The data suggests that the phosphorylation or de-phosphorylation of AMPA receptors is closely associated with the receptor trafficking events. The GluA1 subunit is an important substrate of PP2A, indicating activated PP2A is a critical modulator of AMPA receptor trafficking and might play an important role in AMPA receptor-mediated nociception (Figure 2). In future studies, it will be valuable to determine the activity status of PP2A in the spinal cord dorsal horns and the de-phosphorylation as well as trafficking regulation of AMPA receptor GluA1 or GluA2 subunits by PP2A in different animal models of pain.

PP2A regulates synaptic transmission through influencing transcription factors and subsequent chromatin remodeling

Another way in which phosphatases might regulate synaptic transmission on a longer timescale is through the alternation of chromatin remodeling and gene transcription. Calcium-dependent gene transcription has a critical role in both synaptic plasticity and memory formation. The transcription factor cAMP-response-element-binding protein (CREB) has been demonstrated to be involved in synaptic plasticity and long-term memory, and serves a substrate for various phosphatases as well [57]. In cultured hippocampal neurons, PP1 and/or PP2A are observed to be main CREB phosphatases, and although PP2B does not directly de-phosphorylate CREB, it nonetheless has a key role in regulating CREB-mediated transcription process [83]. Various stimuli evoke the transient phosphorylation of CREB protein, but a more sustained phosphorylation seems to be necessary for a CREB-mediated transcription to occur. It has been shown that PP2A may play a critical role in constraining the progression of information from the synapse to the nucleus (Figure 2). It will be important to learn how the activity dependent regulation of PP2A influences the relevant gene expression. PP2A has been shown to form a signaling complex with CaMKIV that regulates CREB phosphorylation, and thus, a CREB-mediated transcription [51]. PP2A holoenzymes may also negatively regulate NF- κ B-mediated transcriptional activities [84]. Some PP2A regulatory subunits, such as PR55 γ and PR55 δ , are inhibitors of JNK and c-Src kinases, which are important to regulate transcription factors.

The important role for multiple protein kinases in regulation of nociception in animal studies suggests its function on nociception-elicited gene expression through

mediation of transcription factors, such as *c-fos* and CREB. Increased phosphorylation of CREB protein through the activation of glutamate receptors and the PKA, PKC, and CaMKII cascades during central sensitization was reported in several animal models of pain [2,6,10,42]. It suggests an intra-cellular connection between activation of transcription factors and molecular mechanisms mediating stimuli-induced nuclear gene activation through several protein kinase pathways. PP2A may exert a negative action on CREB-mediated transcription-dependent central sensitization, since CREB is an important substrate of PP2A. Future investigations need to determine the regulation of PP2A activity during central sensitization through multiple transcription factors, such as *c-fos*, *c-Jun* and NF- κ B.

Previous studies have suggested that post-translational or epi-genetic modification (acetylation, methylation, phosphorylation, etc.) of histones, and subsequently remodeling of chromatin structure, play a critical role in controlling gene transcription and facilitating long-term changes during synaptic plasticity and central sensitization of pain [85-88]. Chromatin structures are regulated in hippocampal and spinal neurons in response to activation of multiple kinase activation. In particular, it has been shown that the histone acetyltransferase activity of CREB binding protein (CBP) is necessary for synaptic plasticity and central sensitization of pain [89]. CREB can be phosphorylated and activated by different kinases and then it recruits the histone acetyltransferase co-activator CBP and its homologue p300. The recruitment of CBP/p300 and changes in the level of histone acetylation are required for gene transcription. Another study from our group reported that the activated JNK signaling pathway was observed to contribute to the regulation of histone remodeling in peripheral sensory neurons following neurotoxic stimulation [90]. Therefore, PP2A activity may regulate the histone-acetylation induced subsequent neuroepigenetic changes and downstream gene transcription through the de-phosphorylation of CREB.

In contrast to histone acetylase, histone deacetylase (HDAC) may remove acetyl groups from lysine residues of histones, and other non-histone proteins, and reverse the action of histone acetylase and block the gene transcription process. PP2A comprises a family of holoenzyme complexes with diverse biological activities, which mainly depend on individual regulatory subunits. The PP2A heterotrimeric complex was formed by the PP2A-A subunit and the catalytic subunit (PP2A-C), while G5PR as a regulatory subunit exhibits phosphatase activity on histone H1. Class II HDACs are key transcriptional regulators whose activities are controlled via phosphorylation-dependent nucleo-cytoplasmic shuttling [91]. PP2A is responsible for de-phosphorylation of class II HDACs triggering nuclear localization and repression

of target genes, whereas phosphorylation triggers cytoplasmic localization leading to activation of target genes [91] (Figure 2). Recent studies have shown that the HDACs have been implicated in spinal nociception in inflammatory pain [92,93]. It is presumable that PP2A may regulate the gene expression in the spinal cord dorsal horns elicited by peripheral noxious stimuli through the de-phosphorylation of class II HDACs and subsequent neuro-epigenetic alternations.

Concluding remark

In this review, we have highlighted the roles of PP2A in synaptic plasticity and central sensitization of nociceptive process. Specifically, they have key roles in limiting LTP induction and maintenance, and in triggering LTD induction. PP2A activity is regulated by holoenzyme composition, post-translational modification of methylation and phosphorylation. PP2A may exert its action through the de-phosphorylation of critical substrates relevant to nociceptive processing, such as NMDA receptor subunits, AMPA receptor GluA1 subunits, transcription factors and class II HDACs. All these substrate molecules are implicated in synaptic plasticity and central sensitization in the central nervous system. PP2A may serve as a potential molecular target that can be selectively and effectively modulated through pharmaceutical intervention to treat pain.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

YW participated in the design of the review and drafted the manuscript. YL, JW and YM assisted with the preparation of the manuscript and figures. LF, and XZ conceived of the review and participated in its design and helped to draft the manuscript. All authors read and approved the final manuscript.

Acknowledgments

This work was supported by National Natural Science Foundation of China (30801073/H0903, 81171055/H0903), Natural Science Foundation of Beijing (7112054) and New Century Excellent Talents Program from Ministry of Education, China (NCET-10-0014).

Author details

¹Department of Anesthesiology, Beijing Chaoyang Hospital, Capital Medical University, Beijing 100020, China. ²Department of Neuroscience and Cell Biology, The University of Texas Medical Branch, Galveston, TX 77555-0517, USA. ³Department of Neurosurgery, Cancer Center, Sun Yat-sen University, Guangzhou 510060, China. ⁴Department of Neurosurgery and Neurology, Lineberger Cancer Center, School of Medicine, University of North Carolina, Chapel Hill, NC 27599, USA.

Received: 23 February 2013 Accepted: 30 August 2013

Published: 8 September 2013

References

1. Fang L, Wu J, Lin Q, Willis WD: Calcium-calmodulin-dependent protein kinase II contributes to spinal cord central sensitization. *J Neurosci* 2002, **22**:4196–4204.
2. Wu J, Su G, Ma L, Zhang X, Lei Y, Li J, Lin Q, Fang L: Protein kinases mediate increment of the phosphorylation of cyclic AMP-responsive element binding protein in spinal cord of rats following capsaicin injection. *Mol Pain* 2005, **1**:26.
3. Wu J, Li J, Lin Q, Fang L: Signal transduction in chronic pain. *Int Anesthesiol Clin* 2007, **45**:73–81.
4. Faure C, Ramos M, Girault JA: Pyk2 cytonuclear localization: mechanisms and regulation by serine dephosphorylation. *Cell Mol Life Sci* 2013, **70**:137–152.
5. Mizuno K: Signaling mechanisms and functional roles of cofilin phosphorylation and dephosphorylation. *Cell Signal* 2013, **25**:457–469.
6. Fang L, Wu J, Zhang X, Lin Q, Willis WD: Calcium/calmodulin dependent protein kinase II regulates the phosphorylation of cyclic AMP-responsive element-binding protein of spinal cord in rats following noxious stimulation. *Neurosci Lett* 2005, **374**:1–4.
7. Fang L, Wu J, Lin Q, Willis WD: Protein kinases regulate the phosphorylation of the GluR1 subunit of AMPA receptors of spinal cord in rats following noxious stimulation. *Brain Res Mol Brain Res* 2003, **118**:160–165.
8. Wang Y, Wu J, Wu Z, Lin Q, Yue Y, Fang L: Regulation of AMPA receptors in spinal nociception. *Mol Pain* 2010, **6**:5.
9. Wang Y, Mu X, Wu J, Wu A, Fang L, Li J, Yue Y: Differential roles of phosphorylated AMPA receptor GluR1 subunits at Serine-831 and Serine-845 sites in spinal cord dorsal horn in a rat model of post-operative pain. *Neurochem Res* 2011, **36**:170–176.
10. Wu J, Fang L, Lin Q, Willis WD: The role of nitric oxide in the phosphorylation of cyclic adenosine monophosphate-responsive element-binding protein in the spinal cord after intradermal injection of capsaicin. *J Pain* 2002, **3**:190–198.
11. Zhang X, Wu J, Fang L, Willis WD: The effects of protein phosphatase inhibitors on the duration of central sensitization of rat dorsal horn neurons following injection of capsaicin. *Mol Pain* 2006, **2**:23.
12. Zhang X, Wu J, Lei Y, Fang L, Willis WD: Protein phosphatase 2A regulates central sensitization in the spinal cord of rats following intradermal injection of capsaicin. *Mol Pain* 2006, **2**:9.
13. Zhang X, Wu J, Lei Y, Fang L, Willis WD: Protein phosphatase modulates the phosphorylation of spinal cord NMDA receptors in rats following intradermal injection of capsaicin. *Brain Res Mol Brain Res* 2005, **138**:264–272.
14. Zhang X, Wu J, Fang L, Willis WD: The effects of protein phosphatase inhibitors on nociceptive behavioral responses of rats following intradermal injection of capsaicin. *Pain* 2003, **106**:443–451.
15. Gabra BH, Bailey CP, Kelly E, Sanders AV, Henderson G, Smith FL, Dewey WL: Evidence for an important role of protein phosphatases in the mechanism of morphine tolerance. *Brain Res* 2007, **1159**:86–93.
16. Zhang Q, Claret FX: Phosphatases: the new brakes for cancer development? *Enzyme Res* 2012, **2012**:659649.
17. Winder DG, Sweatt JD: Roles of serine/threonine phosphatases in hippocampal synaptic plasticity. *Nat Rev Neurosci* 2001, **2**:461–474.
18. Janssens V, Goris J: Protein phosphatase 2A: a highly regulated family of serine/threonine phosphatases implicated in cell growth and signalling. *Biochem J* 2001, **353**:417–439.
19. Janssens V, Longin S, Goris J: PP2A holoenzyme assembly: in cauda venenum (the sting is in the tail). *Trends Biochem Sci* 2008, **33**:113–121.
20. Janssens V, Derua R, Zwaenepoel K, Waelkens E, Goris J: Specific regulation of protein phosphatase 2A PR72/B" subunits by calpain. *Biochem Biophys Res Commun* 2009, **386**:676–681.
21. Park JH, Sung HY, Lee JY, Kim HJ, Ahn JH, Jo I: B56alpha subunit of protein phosphatase 2A mediates retinoic acid-induced decreases in phosphorylation of endothelial nitric oxide synthase at serine 1179 and nitric oxide production in bovine aortic endothelial cells. *Biochem Biophys Res Commun* 2013, **430**:476–481.
22. Zwaenepoel K, Louis JV, Goris J, Janssens V: Diversity in genomic organisation, developmental regulation and distribution of the murine PR72/B" subunits of protein phosphatase 2A. *BMC Genomics* 2008, **9**:393.
23. Ahn JH, Sung JY, McAvoy T, Nishi A, Janssens V, Goris J, Greengard P, Naim AC: The B"PR72 subunit mediates Ca2+-dependent dephosphorylation of DARPP-32 by protein phosphatase 2A. *Proc Natl Acad Sci U S A* 2007, **104**:9876–9881.
24. Lechward K, Awotunde OS, Swiatek W, Muszynska G: Protein phosphatase 2A: variety of forms and diversity of functions. *Acta Biochim Pol* 2001, **48**:921–933.
25. Cho US, Morrone S, Sablina AA, Arroyo JD, Hahn WC, Xu W: Structural basis of PP2A inhibition by small t antigen. *PLoS Biol* 2007, **5**:e202.
26. Letourneux C, Rocher G, Porteu F: B56-containing PP2A dephosphorylate ERK and their activity is controlled by the early gene IEX-1 and ERK. *EMBO J* 2006, **25**:727–738.

27. Chen J, Martin BL, Brautigam DL: Regulation of protein serine-threonine phosphatase type-2A by tyrosine phosphorylation. *Science* 1992, **257**:1261–1264.
28. Longin S, Zwaenepoel K, Martens E, Louis JV, Rondelez E, Goris J, Janssens V: Spatial control of protein phosphatase 2A (de)methylation. *Exp Cell Res* 2008, **314**:68–81.
29. Longin S, Zwaenepoel K, Louis JV, Dilworth S, Goris J, Janssens V: Selection of protein phosphatase 2A regulatory subunits is mediated by the C terminus of the catalytic subunit. *J Biol Chem* 2007, **282**:26971–26980.
30. Ogris E, Gibson DM, Pallas DC: Protein phosphatase 2A subunit assembly: the catalytic subunit carboxyl terminus is important for binding cellular B subunit but not polyomavirus middle tumor antigen. *Oncogene* 1997, **15**:911–917.
31. Bryant JC, Westphal RS, Wadzinski BE: Methylated C-terminal leucine residue of PP2A catalytic subunit is important for binding of regulatory Balpha subunit. *Biochem J* 1999, **339**(Pt 2):241–246.
32. Leulliot N, Quevillon-Cheruel S, Sorel I, de La Sierra-Gallay L, Collinet B, Graille M, Blondeau K, Bettache N, Poupon A, Janin J, van TH: Structure of protein phosphatase methyltransferase 1 (PPM1), a leucine carboxyl methyltransferase involved in the regulation of protein phosphatase 2A activity. *J Biol Chem* 2004, **279**:8351–8358.
33. Rudrabhatla P, Pant HC: Role of protein phosphatase 2A in Alzheimer's disease. *Curr Alzheimer Res* 2011, **8**:623–632.
34. Torrent L, Ferrer I: PP2A and Alzheimer disease. *Curr Alzheimer Res* 2012, **9**:248–256.
35. Sun XY, Wei YP, Xiong Y, Wang XC, Xie AJ, Wang XL, Yang Y, Wang Q, Lu YM, Liu R, Wang JZ: Synaptic released zinc promotes tau hyperphosphorylation by inhibition of protein phosphatase 2A (PP2A). *J Biol Chem* 2012, **287**:11174–11182.
36. Sontag E, Hladik C, Montgomery L, Luangpirom A, Mudrak I, Ogris E, White CL III: Downregulation of protein phosphatase 2A carboxyl methylation and methyltransferase may contribute to Alzheimer disease pathogenesis. *J Neuropathol Exp Neurol* 2004, **63**:1080–1091.
37. Sontag E, Luangpirom A, Hladik C, Mudrak I, Ogris E, Speciale S, White CL III: Altered expression levels of the protein phosphatase 2A ABalpaC enzyme are associated with Alzheimer disease pathology. *J Neuropathol Exp Neurol* 2004, **63**:287–301.
38. Sandkuhler J, Gruber-Schoffnegger D: Hyperalgesia by synaptic long-term potentiation (LTP): an update. *Curr Opin Pharmacol* 2012, **12**:18–27.
39. Sandkuhler J: Central sensitization versus synaptic long-term potentiation (LTP): a critical comment. *J Pain* 2010, **11**:798–800.
40. Larsson M, Broman J: Synaptic plasticity and pain: role of ionotropic glutamate receptors. *Neuroscientist* 2011, **17**:256–273.
41. Larsson M, Broman J: Translocation of GluR1-containing AMPA receptors to a spinal nociceptive synapse during acute noxious stimulation. *J Neurosci* 2008, **28**:7084–7090.
42. Wu J, Su G, Ma L, Zhang X, Lei Y, Lin Q, Nauta HJ, Li J, Fang L: The role of c-AMP-dependent protein kinase in spinal cord and post synaptic dorsal column neurons in a rat model of visceral pain. *Neurochem Int* 2007, **50**:710–718.
43. Willis WD: Role of neurotransmitters in sensitization of pain responses. *Ann N Y Acad Sci* 2001, **933**:142–156.
44. Tanabe M, Nagatani Y, Saitoh K, Takasu K, Ono H: Pharmacological assessments of nitric oxide synthase isoforms and downstream diversity of NO signaling in the maintenance of thermal and mechanical hypersensitivity after peripheral nerve injury in mice. *Neuropharmacology* 2009, **56**:702–708.
45. Schmidtke A, Ruth P, Geisslinger G, Tegeder I: Inhibition of cyclic guanosine 5'-monophosphate-dependent protein kinase I (PKG-I) in lumbar spinal cord reduces formalin-induced hyperalgesia and PKG upregulation. *Nitric Oxide* 2003, **8**:89–94.
46. Bliss TV, Collingridge GL: A synaptic model of memory: long-term potentiation in the hippocampus. *Nature* 1993, **361**:31–39.
47. Thiels E, Norman ED, Barrionuevo G, Klann E: Transient and persistent increases in protein phosphatase activity during long-term depression in the adult hippocampus in vivo. *Neuroscience* 1998, **86**:1023–1029.
48. Figurov A, Boddeke H, Muller D: Enhancement of AMPA-mediated synaptic transmission by the protein phosphatase inhibitor calyculin A in rat hippocampal slices. *Eur J Neurosci* 1993, **5**:1035–1041.
49. Blitzer RD, Wong T, Nouranifar R, Iyengar R, Landau EM: Postsynaptic cAMP pathway gates early LTP in hippocampal CA1 region. *Neuron* 1995, **15**:1403–1414.
50. Winder DG, Mansuy IM, Osman M, Moallem TM, Kandel ER: Genetic and pharmacological evidence for a novel, intermediate phase of long-term potentiation suppressed by calcineurin. *Cell* 1998, **92**:25–37.
51. Blitzer RD, Connor JH, Brown GP, Wong T, Shenolikar S, Iyengar R, Landau EM: Gating of CaMKII by cAMP-regulated protein phosphatase activity during LTP. *Science* 1998, **280**:1940–1942.
52. Makhinson M, Chotiner JK, Watson JB, O'Dell TJ: Adenylyl cyclase activation modulates activity-dependent changes in synaptic strength and Ca²⁺/calmodulin-dependent kinase II autophosphorylation. *J Neurosci* 1999, **19**:2500–2510.
53. Fukunaga K, Muller D, Ohmitsu M, Bako E, DePaoli-Roach AA, Miyamoto E: Decreased protein phosphatase 2A activity in hippocampal long-term potentiation. *J Neurochem* 2000, **74**:807–817.
54. Strack S, Choi S, Lovinger DM, Colbran RJ: Translocation of autophosphorylated calcium/calmodulin-dependent protein kinase II to the postsynaptic density. *J Biol Chem* 1997, **272**:13467–13470.
55. Pi HJ, Lisman JE: Coupled phosphatase and kinase switches produce the tristability required for long-term potentiation and long-term depression. *J Neurosci* 2008, **28**:13132–13138.
56. Mulkey RM, Herron CE, Malenka RC: An essential role for protein phosphatases in hippocampal long-term depression. *Science* 1993, **261**:1051–1055.
57. Mauna JC, Miyamae T, Pulli B, Thiels E: Protein phosphatases 1 and 2A are both required for long-term depression and associated dephosphorylation of cAMP response element binding protein in hippocampal area CA1 in vivo. *Hippocampus* 2011, **21**:1093–1104.
58. Woo NH, Nguyen PV: "Silent" metaplasticity of the late phase of long-term potentiation requires protein phosphatases. *Learn Mem* 2002, **9**:202–213.
59. Jouvenceau A, Billard JM, Haditsch U, Mansuy IM, Dutar P: Different phosphatase-dependent mechanisms mediate long-term depression and depotentiation of long-term potentiation in mouse hippocampal CA1 area. *Eur J Neurosci* 2003, **18**:1279–1285.
60. Tapia R, Pena F, Arias C: Neurotoxic and synaptic effects of okadaic acid, an inhibitor of protein phosphatases. *Neurochem Res* 1999, **24**:1423–1430.
61. Huang CC, Liang YC, Hsu KS: Characterization of the mechanism underlying the reversal of long term potentiation by low frequency stimulation at hippocampal CA1 synapses. *J Biol Chem* 2001, **276**:48108–48117.
62. Arias C, Montiel T, Pena F, Ferrera P, Tapia R: Okadaic acid induces epileptic seizures and hyperphosphorylation of the NR2B subunit of the NMDA receptor in rat hippocampus in vivo. *Exp Neurol* 2002, **177**:284–291.
63. Chan SF, Sucher NJ: An NMDA receptor signaling complex with protein phosphatase 2A. *J Neurosci* 2001, **21**:7985–7992.
64. Atianjoh FE, Yaster M, Zhao X, Takamiya K, Xia J, Gauda EB, Haganir RL, Tao YX: Spinal cord protein interacting with C kinase 1 is required for the maintenance of complete Freund's adjuvant-induced inflammatory pain but not for incision-induced post-operative pain. *Pain* 2010, **151**:226–234.
65. Kristensen AS, Jenkins MA, Banke TG, Schousboe A, Makino Y, Johnson RC, Haganir R, Traynelis SF: Mechanism of Ca²⁺/calmodulin-dependent kinase II regulation of AMPA receptor gating. *Nat Neurosci* 2011, **14**:727–735.
66. Anggono V, Haganir RL: Regulation of AMPA receptor trafficking and synaptic plasticity. *Curr Opin Neurobiol* 2012, **22**:461–469.
67. Park JS, Voitenko N, Petralia RS, Guan X, Xu JT, Steinberg JP, Takamiya K, Sotnik A, Kopach O, Haganir RL, Tao YX: Persistent inflammation induces GluR2 internalization via NMDA receptor-triggered PKC activation in dorsal horn neurons. *J Neurosci* 2009, **29**:3206–3219.
68. Clem RL, Anggono V, Haganir RL: PICK1 regulates incorporation of calcium-permeable AMPA receptors during cortical synaptic strengthening. *J Neurosci* 2010, **30**:6360–6366.
69. Lee HK, Kameyama K, Haganir RL, Bear MF: NMDA induces long-term synaptic depression and dephosphorylation of the GluR1 subunit of AMPA receptors in hippocampus. *Neuron* 1998, **21**:1151–1162.
70. Kameyama K, Lee HK, Bear MF, Haganir RL: Involvement of a postsynaptic protein kinase A substrate in the expression of homosynaptic long-term depression. *Neuron* 1998, **21**:1163–1175.
71. Larsson M, Broman J: Pathway-specific bidirectional regulation of Ca²⁺/calmodulin-dependent protein kinase II at spinal nociceptive synapses after acute noxious stimulation. *J Neurosci* 2006, **26**:4198–4205.
72. Lee HK, Barbarosie M, Kameyama K, Bear MF, Haganir RL: Regulation of distinct AMPA receptor phosphorylation sites during bidirectional synaptic plasticity. *Nature* 2000, **405**:955–959.

73. Lee HK, Takamiya K, Han JS, Man H, Kim CH, Rumbaugh G, Yu S, Ding L, He C, Petralia RS, Wenthold RJ, Gallagher M, Huganir RL: **Phosphorylation of the AMPA receptor GluR1 subunit is required for synaptic plasticity and retention of spatial memory.** *Cell* 2003, **112**:631–643.
74. Lu W, Roche KW: **Posttranslational regulation of AMPA receptor trafficking and function.** *Curr Opin Neurobiol* 2012, **22**:470–479.
75. Peng HY, Chen GD, Hsieh MC, Lai CY, Huang YP, Lin TB: **Spinal SGK1/GRASP-1/Rab4 is involved in complete Freund's adjuvant-induced inflammatory pain via regulating dorsal horn GluR1-containing AMPA receptor trafficking in rats.** *Pain* 2012, **153**:2380–2392.
76. Beattie EC, Carroll RC, Yu X, Morishita W, Yasuda H, von ZM, Malenka RC: **Regulation of AMPA receptor endocytosis by a signaling mechanism shared with LTD.** *Nat Neurosci* 2000, **3**:1291–1300.
77. Clem RL, Huganir RL: **Calcium-permeable AMPA receptor dynamics mediate fear memory erasure.** *Science* 2010, **330**:1108–1112.
78. Esteban JA, Shi SH, Wilson C, Nuriya M, Huganir RL, Malinow R: **PKA phosphorylation of AMPA receptor subunits controls synaptic trafficking underlying plasticity.** *Nat Neurosci* 2003, **6**:136–143.
79. Man HY, Sekine-Aizawa Y, Huganir RL: **Regulation of [alpha]-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor trafficking through PKA phosphorylation of the Glu receptor 1 subunit.** *Proc Natl Acad Sci USA* 2007, **104**:3579–3584.
80. Hayashi Y, Shi SH, Esteban JA, Piccini A, Poncer JC, Malinow R: **Driving AMPA receptors into synapses by LTP and CaMKII: requirement for GluR1 and PDZ domain interaction.** *Science* 2000, **287**:2262–2267.
81. Tao YX: **AMPA receptor trafficking in inflammation-induced dorsal horn central sensitization.** *Neurosci Bull* 2012, **28**:111–120.
82. Choi JJ, Svensson CI, Koehn FJ, Bhushkute A, Sorokin LS: **Peripheral inflammation induces tumor necrosis factor dependent AMPA receptor trafficking and Akt phosphorylation in spinal cord in addition to pain behavior.** *Pain* 2010, **149**:243–253.
83. Huang L, Balsara RD, Sheng Z, Castellino FJ: **Conantokins inhibit NMDAR-dependent calcium influx in developing rat hippocampal neurons in primary culture with resulting effects on CREB phosphorylation.** *Mol Cell Neurosci* 2010, **45**:163–172.
84. Hsieh CY, Hsu MJ, Hsiao G, Wang YH, Huang CW, Chen SW, Jayakumar T, Chiu PT, Chiu YH, Sheu JR: **Andrographolide enhances nuclear factor-kappaB subunit p65 Ser536 dephosphorylation through activation of protein phosphatase 2A in vascular smooth muscle cells.** *J Biol Chem* 2011, **286**:5942–5955.
85. Denk F, McMahon SB: **Chronic pain: emerging evidence for the involvement of epigenetics.** *Neuron* 2012, **73**:435–444.
86. Sultan FA, Day JJ: **Epigenetic mechanisms in memory and synaptic function.** *Epigenomics* 2011, **3**:157–181.
87. Geranton SM: **Targeting epigenetic mechanisms for pain relief.** *Curr Opin Pharmacol* 2012, **12**:35–41.
88. Wang Y, Chen Z, Zhao Y, Shi R, Wang Y, Xu J, Wu A, Johns RA, Yue Y: **Epigenetics as a new therapeutic target for postoperative cognitive dysfunction.** *Med Hypotheses* 2013, **80**:249–251.
89. Zhu XY, Huang CS, Li Q, Chang RM, Song ZB, Zou WY, Guo QL: **p300 exerts an epigenetic role in chronic neuropathic pain through its acetyltransferase activity in rats following chronic constriction injury (CCI).** *Mol Pain* 2012, **8**:84.
90. Wu J, Zhang X, Nauta HJ, Lin Q, Li J, Fang L: **JNK1 regulates histone acetylation in trigeminal neurons following chemical stimulation.** *Biochem Biophys Res Commun* 2008, **376**:781–786.
91. Paroni G, Cernotta N, Dello RC, Gallinari P, Pallaoro M, Foti C, Talamo F, Orsatti L, Steinkuhler C, Brancolini C: **PP2A regulates HDAC4 nuclear import.** *Mol Biol Cell* 2008, **19**:655–667.
92. Tochiki KK, Cunningham J, Hunt SP, Geranton SM: **The expression of spinal methyl-CpG-binding protein 2, DNA methyltransferases and histone deacetylases is modulated in persistent pain states.** *Mol Pain* 2012, **8**:14.
93. Zhang Z, Cai YQ, Zou F, Bie B, Pan ZZ: **Epigenetic suppression of GAD65 expression mediates persistent pain.** *Nat Med* 2011, **17**:1448–1455.

doi:10.1186/1744-8069-9-46

Cite this article as: Wang et al.: Roles of phosphatase 2A in nociceptive signal processing. *Molecular Pain* 2013 **9**:46.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

